

GEORGIAN MEDICAL NEWS

ISSN 1512-0112

№ 4 (313) Апрель 2021

ТБИЛИСИ - NEW YORK



ЕЖЕМЕСЯЧНЫЙ НАУЧНЫЙ ЖУРНАЛ

Медицинские новости Грузии
cfnfhsdtkj c cfvtl bwbyj cbf[k tyb

GEORGIAN MEDICAL NEWS

No 4 (313) 2021

Published in cooperation with and under the patronage
of the Tbilisi State Medical University

Издается в сотрудничестве и под патронажем
Тбилисского государственного медицинского университета

გამოიცემა თბილისის სახელმწიფო სამედიცინო უნივერსიტეტთან
თანამშრომლობითა და მისი პატრონაჟით

ЕЖЕМЕСЯЧНЫЙ НАУЧНЫЙ ЖУРНАЛ
ТБИЛИСИ - НЬЮ-ЙОРК

GMN: Georgian Medical News is peer-reviewed, published monthly journal committed to promoting the science and art of medicine and the betterment of public health, published by the GMN Editorial Board and The International Academy of Sciences, Education, Industry and Arts (U.S.A.) since 1994. **GMN** carries original scientific articles on medicine, biology and pharmacy, which are of experimental, theoretical and practical character; publishes original research, reviews, commentaries, editorials, essays, medical news, and correspondence in English and Russian.

GMN is indexed in MEDLINE, SCOPUS, PubMed and VINITI Russian Academy of Sciences. The full text content is available through EBSCO databases.

GMN: Медицинские новости Грузии - ежемесячный рецензируемый научный журнал, издаётся Редакционной коллегией и Международной академией наук, образования, искусств и естествознания (IASEIA) США с 1994 года на русском и английском языках в целях поддержки медицинской науки и улучшения здравоохранения. В журнале публикуются оригинальные научные статьи в области медицины, биологии и фармации, статьи обзорного характера, научные сообщения, новости медицины и здравоохранения.

Журнал индексируется в MEDLINE, отражён в базе данных SCOPUS, PubMed и ВИНТИ РАН. Полнотекстовые статьи журнала доступны через БД EBSCO.

GMN: Georgian Medical News – საქართველოს სამედიცინო სიახლენი – არის ყოველთვიური სამეცნიერო სამედიცინო რეცენზირებადი ჟურნალი, გამოიცემა 1994 წლიდან, წარმოადგენს სარედაქციო კოლეგიისა და აშშ-ის მეცნიერების, განათლების, ინდუსტრიის, ხელოვნებისა და ბუნებისმეტყველების საერთაშორისო აკადემიის ერთობლივ გამოცემას. GMN-ში რუსულ და ინგლისურ ენებზე ქვეყნდება ექსპერიმენტული, თეორიული და პრაქტიკული ხასიათის ორიგინალური სამეცნიერო სტატიები მედიცინის, ბიოლოგიისა და ფარმაციის სფეროში, მიმოხილვითი ხასიათის სტატიები.

ჟურნალი ინდექსირებულია MEDLINE-ის საერთაშორისო სისტემაში, ასახულია SCOPUS-ის, PubMed-ის და ВИНТИ РАН-ის მონაცემთა ბაზებში. სტატიების სრული ტექსტი ხელმისაწვდომია EBSCO-ს მონაცემთა ბაზებშიდან.

МЕДИЦИНСКИЕ НОВОСТИ ГРУЗИИ

Ежемесячный совместный грузино-американский научный электронно-печатный журнал
Агентства медицинской информации Ассоциации деловой прессы Грузии,
Международной академии наук, индустрии, образования и искусств США.
Издается с 1994 г., распространяется в СНГ, ЕС и США

ГЛАВНЫЙ РЕДАКТОР

Николай Пирцхалаишвили

НАУЧНЫЙ РЕДАКТОР

Елене Гиоргадзе

ЗАМЕСТИТЕЛЬ ГЛАВНОГО РЕДАКТОРА

Нино Микаберидзе

НАУЧНО-РЕДАКЦИОННЫЙ СОВЕТ

Зураб Вадачкориа - председатель Научно-редакционного совета

Михаил Бахмутский (США), Александр Геннинг (Германия), Амиран Гамкрелидзе (Грузия),
Константин Кипиани (Грузия), Георгий Камкамидзе (Грузия),
Паата Куртанидзе (Грузия), Вахтанг Масхулия (Грузия),
Тенгиз Ризнис (США), Реваз Сепиашвили (Грузия), Дэвид Элуа (США)

НАУЧНО-РЕДАКЦИОННАЯ КОЛЛЕГИЯ

Константин Кипиани - председатель Научно-редакционной коллегии

Архимандрит Адам - Вахтанг Ахаладзе, Амиран Антадзе, Нелли Антелава, Тенгиз Асатиани,
Гия Берадзе, Рима Бериашвили, Лео Бокерия, Отар Герзмава, Лиана Гогиашвили, Нодар Гогебашвили,
Николай Гонгадзе, Лия Дваладзе, Тамар Долиашвили, Манана Жвания, Тамар Зерекидзе,
Ирина Квачадзе, Нана Квирквелия, Зураб Кеванишвили, Гурам Кикнадзе, Димитрий
Кордзаиа, Теймураз Лежава, Нодар Ломидзе, Джанлуиджи Мелотти, Марина Мамаладзе,
Караман Пагава, Мамука Пирцхалаишвили, Анна Рехвиашвили, Мака Сологашвили, Рамаз Хецуриани,
Рудольф Хохенфеллнер, Кахабер Челидзе, Тинатин Чиковани, Арчил Чхотуа,
Рамаз Шенгелия, Кетеван Эбралидзе

Website:

www.geomednews.org

The International Academy of Sciences, Education, Industry & Arts. P.O.Box 390177,
Mountain View, CA, 94039-0177, USA. Tel/Fax: (650) 967-4733

Версия: печатная. **Цена:** свободная.

Условия подписки: подписка принимается на 6 и 12 месяцев.

По вопросам подписки обращаться по тел.: 293 66 78.

Контактный адрес: Грузия, 0177, Тбилиси, ул. Асатиани 7, IV этаж, комната 408
тел.: 995(32) 254 24 91, 5(55) 75 65 99

Fax: +995(32) 253 70 58, e-mail: ninomikaber@geomednews.com; nikopir@geomednews.com

По вопросам размещения рекламы обращаться по тел.: 5(99) 97 95 93

© 2001. Ассоциация деловой прессы Грузии

© 2001. The International Academy of Sciences,
Education, Industry & Arts (USA)

GEORGIAN MEDICAL NEWS

Monthly Georgia-US joint scientific journal published both in electronic and paper formats of the Agency of Medical Information of the Georgian Association of Business Press; International Academy of Sciences, Education, Industry and Arts (USA).
Published since 1994. Distributed in NIS, EU and USA.

EDITOR IN CHIEF

Nicholas Pirtskhalaishvili

SCIENTIFIC EDITOR

Elene Giorgadze

DEPUTY CHIEF EDITOR

Nino Mikaberidze

SCIENTIFIC EDITORIAL COUNCIL

Zurab Vadachkoria - Head of Editorial council

Michael Bakhmutsky (USA), Alexander Gënning (Germany),
Amiran Gamkrelidze (Georgia), David Elua (USA),
Konstantin Kipiani (Georgia), Giorgi Kamkamidze (Georgia), Paata Kurtanidze (Georgia),
Vakhtang Maskhulia (Georgia), Tengiz Riznis (USA), Revaz Sepiashvili (Georgia)

SCIENTIFIC EDITORIAL BOARD

Konstantin Kipiani - Head of Editorial board

Archimandrite Adam - Vakhtang Akhaladze, Amiran Antadze, Nelly Antelava,
Tengiz Asatiani, Gia Beradze, Rima Beriashvili, Leo Bokeria, Kakhaber Chelidze,
Tinatin Chikovani, Archil Chkhotua, Lia Dvaladze, Tamar Doliashvili, Ketevan Ebralidze,
Otar Gerzmava, Liana Gogiashvili, Nodar Gogebashvili, Nicholas Gongadze,
Rudolf Hohenfellner, Zurab Kevanishvili, Ramaz Khetsuriani, Guram Kiknadze,
Dimitri Kordzaia, Irina Kvachadze, Nana Kvirkvelia, Teymuraz Lezhava, Nodar Lomidze, Marina
Mamaladze, Gianluigi Melotti, Kharaman Pagava, Mamuka Pirtskhalaishvili,
Anna Rekhviashvili, Maka Sologhashvili, Ramaz Shengelia, Tamar Zerekidze, Manana Zhvania

CONTACT ADDRESS IN TBILISI

GMN Editorial Board
7 Asatiani Street, 4th Floor
Tbilisi, Georgia 0177

Phone: 995 (32) 254-24-91
995 (32) 253-70-58
Fax: 995 (32) 253-70-58

CONTACT ADDRESS IN NEW YORK

NINITEX INTERNATIONAL, INC.
3 PINE DRIVE SOUTH
ROSLYN, NY 11576 U.S.A.

Phone: +1 (917) 327-7732

WEBSITE

www.geomednews.org

К СВЕДЕНИЮ АВТОРОВ!

При направлении статьи в редакцию необходимо соблюдать следующие правила:

1. Статья должна быть представлена в двух экземплярах, на русском или английском языках, напечатанная через **полтора интервала на одной стороне стандартного листа с шириной левого поля в три сантиметра**. Используемый компьютерный шрифт для текста на русском и английском языках - **Times New Roman (Кириллица)**, для текста на грузинском языке следует использовать **AcadNusx**. Размер шрифта - **12**. К рукописи, напечатанной на компьютере, должен быть приложен CD со статьей.

2. Размер статьи должен быть не менее десяти и не более двадцати страниц машинописи, включая указатель литературы и резюме на английском, русском и грузинском языках.

3. В статье должны быть освещены актуальность данного материала, методы и результаты исследования и их обсуждение.

При представлении в печать научных экспериментальных работ авторы должны указывать вид и количество экспериментальных животных, применявшиеся методы обезболивания и усыпления (в ходе острых опытов).

4. К статье должны быть приложены краткое (на полстраницы) резюме на английском, русском и грузинском языках (включающее следующие разделы: цель исследования, материал и методы, результаты и заключение) и список ключевых слов (key words).

5. Таблицы необходимо представлять в печатной форме. Фотокопии не принимаются. **Все цифровые, итоговые и процентные данные в таблицах должны соответствовать таковым в тексте статьи**. Таблицы и графики должны быть озаглавлены.

6. Фотографии должны быть контрастными, фотокопии с рентгенограмм - в позитивном изображении. Рисунки, чертежи и диаграммы следует озаглавить, пронумеровать и вставить в соответствующее место текста **в tiff формате**.

В подписях к микрофотографиям следует указывать степень увеличения через окуляр или объектив и метод окраски или импрегнации срезов.

7. Фамилии отечественных авторов приводятся в оригинальной транскрипции.

8. При оформлении и направлении статей в журнал МНГ просим авторов соблюдать правила, изложенные в «Единых требованиях к рукописям, представляемым в биомедицинские журналы», принятых Международным комитетом редакторов медицинских журналов - <http://www.spinesurgery.ru/files/publish.pdf> и http://www.nlm.nih.gov/bsd/uniform_requirements.html В конце каждой оригинальной статьи приводится библиографический список. В список литературы включаются все материалы, на которые имеются ссылки в тексте. Список составляется в алфавитном порядке и нумеруется. Литературный источник приводится на языке оригинала. В списке литературы сначала приводятся работы, написанные знаками грузинского алфавита, затем кириллицей и латиницей. Ссылки на цитируемые работы в тексте статьи даются в квадратных скобках в виде номера, соответствующего номеру данной работы в списке литературы. Большинство цитированных источников должны быть за последние 5-7 лет.

9. Для получения права на публикацию статья должна иметь от руководителя работы или учреждения визу и сопроводительное отношение, написанные или напечатанные на бланке и заверенные подписью и печатью.

10. В конце статьи должны быть подписи всех авторов, полностью приведены их фамилии, имена и отчества, указаны служебный и домашний номера телефонов и адреса или иные координаты. Количество авторов (соавторов) не должно превышать пяти человек.

11. Редакция оставляет за собой право сокращать и исправлять статьи. Корректур авторам не высылаются, вся работа и сверка проводится по авторскому оригиналу.

12. Недопустимо направление в редакцию работ, представленных к печати в иных издательствах или опубликованных в других изданиях.

При нарушении указанных правил статьи не рассматриваются.

REQUIREMENTS

Please note, materials submitted to the Editorial Office Staff are supposed to meet the following requirements:

1. Articles must be provided with a double copy, in English or Russian languages and typed or computer-printed on a single side of standard typing paper, with the left margin of 3 centimeters width, and 1.5 spacing between the lines, typeface - **Times New Roman (Cyrillic)**, print size - 12 (referring to Georgian and Russian materials). With computer-printed texts please enclose a CD carrying the same file titled with Latin symbols.

2. Size of the article, including index and resume in English, Russian and Georgian languages must be at least 10 pages and not exceed the limit of 20 pages of typed or computer-printed text.

3. Submitted material must include a coverage of a topical subject, research methods, results, and review.

Authors of the scientific-research works must indicate the number of experimental biological species drawn in, list the employed methods of anesthetization and soporific means used during acute tests.

4. Articles must have a short (half page) abstract in English, Russian and Georgian (including the following sections: aim of study, material and methods, results and conclusions) and a list of key words.

5. Tables must be presented in an original typed or computer-printed form, instead of a photocopied version. **Numbers, totals, percentile data on the tables must coincide with those in the texts of the articles.** Tables and graphs must be headed.

6. Photographs are required to be contrasted and must be submitted with doubles. Please number each photograph with a pencil on its back, indicate author's name, title of the article (short version), and mark out its top and bottom parts. Drawings must be accurate, drafts and diagrams drawn in Indian ink (or black ink). Photocopies of the X-ray photographs must be presented in a positive image in **tiff format**.

Accurately numbered subtitles for each illustration must be listed on a separate sheet of paper. In the subtitles for the microphotographs please indicate the ocular and objective lens magnification power, method of coloring or impregnation of the microscopic sections (preparations).

7. Please indicate last names, first and middle initials of the native authors, present names and initials of the foreign authors in the transcription of the original language, enclose in parenthesis corresponding number under which the author is listed in the reference materials.

8. Please follow guidance offered to authors by The International Committee of Medical Journal Editors guidance in its Uniform Requirements for Manuscripts Submitted to Biomedical Journals publication available online at: http://www.nlm.nih.gov/bsd/uniform_requirements.html
http://www.icmje.org/urm_full.pdf

In GMN style for each work cited in the text, a bibliographic reference is given, and this is located at the end of the article under the title "References". All references cited in the text must be listed. The list of references should be arranged alphabetically and then numbered. References are numbered in the text [numbers in square brackets] and in the reference list and numbers are repeated throughout the text as needed. The bibliographic description is given in the language of publication (citations in Georgian script are followed by Cyrillic and Latin).

9. To obtain the rights of publication articles must be accompanied by a visa from the project instructor or the establishment, where the work has been performed, and a reference letter, both written or typed on a special signed form, certified by a stamp or a seal.

10. Articles must be signed by all of the authors at the end, and they must be provided with a list of full names, office and home phone numbers and addresses or other non-office locations where the authors could be reached. The number of the authors (co-authors) must not exceed the limit of 5 people.

11. Editorial Staff reserves the rights to cut down in size and correct the articles. Proof-sheets are not sent out to the authors. The entire editorial and collation work is performed according to the author's original text.

12. Sending in the works that have already been assigned to the press by other Editorial Staffs or have been printed by other publishers is not permissible.

**Articles that Fail to Meet the Aforementioned
Requirements are not Assigned to be Reviewed.**

ავტორთა საქურაღებოლ!

რედაქციაში სტატიის წარმოდგენისას საჭიროა დაიცვათ შემდეგი წესები:

1. სტატია უნდა წარმოადგინოთ 2 ცალად, რუსულ ან ინგლისურ ენებზე დაბეჭდილი სტანდარტული ფურცლის 1 გვერდზე, 3 სმ სიგანის მარცხენა ველისა და სტრიქონებს შორის 1,5 ინტერვალის დაცვით. გამოყენებული კომპიუტერული შრიფტი რუსულ და ინგლისურენოვან ტექსტებში - **Times New Roman (Кириллица)**, ხოლო ქართულენოვან ტექსტში საჭიროა გამოვიყენოთ **AcadNusx**. შრიფტის ზომა – 12. სტატიას თან უნდა ახლდეს CD სტატიით.

2. სტატიის მოცულობა არ უნდა შეადგენდეს 10 გვერდზე ნაკლებს და 20 გვერდზე მეტს ლიტერატურის სიის და რეზიუმეების (ინგლისურ, რუსულ და ქართულ ენებზე) ჩათვლით.

3. სტატიაში საჭიროა გაშუქდეს: საკითხის აქტუალობა; კვლევის მიზანი; საკვლევი მასალა და გამოყენებული მეთოდები; მიღებული შედეგები და მათი განსჯა. ექსპერიმენტული ხასიათის სტატიების წარმოდგენისას ავტორებმა უნდა მიუთითონ საექსპერიმენტო ცხოველების სახეობა და რაოდენობა; გაუტკივარებისა და დაძინების მეთოდები (მწვავე ცდების პირობებში).

4. სტატიას თან უნდა ახლდეს რეზიუმე ინგლისურ, რუსულ და ქართულ ენებზე არანაკლებ ნახევარი გვერდის მოცულობისა (სათაურის, ავტორების, დაწესებულების მითითებით და უნდა შეიცავდეს შემდეგ განყოფილებებს: მიზანი, მასალა და მეთოდები, შედეგები და დასკვნები; ტექსტუალური ნაწილი არ უნდა იყოს 15 სტრიქონზე ნაკლები) და საკვანძო სიტყვების ჩამონათვალი (key words).

5. ცხრილები საჭიროა წარმოადგინოთ ნაბეჭდი სახით. ყველა ციფრული, შემაჯამებელი და პროცენტული მონაცემები უნდა შეესაბამებოდეს ტექსტში მოყვანილს.

6. ფოტოსურათები უნდა იყოს კონტრასტული; სურათები, ნახაზები, დიაგრამები - დასათაურებული, დანომრილი და სათანადო ადგილას ჩასმული. რენტგენოგრამების ფოტოასლები წარმოადგინეთ პოზიტიური გამოსახულებით **tiff** ფორმატში. მიკროფოტოსურათების წარწერებში საჭიროა მიუთითოთ ოკულარის ან ობიექტივის საშუალებით გადიდების ხარისხი, ანათალების შედეგის ან იმპრეგნაციის მეთოდი და აღნიშნოთ სურათის ზედა და ქვედა ნაწილები.

7. სამამულო ავტორების გვარები სტატიაში აღინიშნება ინიციალების თანდართვით, უცხოურისა – უცხოური ტრანსკრიპციით.

8. სტატიას თან უნდა ახლდეს ავტორის მიერ გამოყენებული სამამულო და უცხოური შრომების ბიბლიოგრაფიული სია (ბოლო 5-8 წლის სიღრმით). ანბანური წყობით წარმოდგენილ ბიბლიოგრაფიულ სიაში მიუთითეთ ჯერ სამამულო, შემდეგ უცხოელი ავტორები (გვარი, ინიციალები, სტატიის სათაური, ჟურნალის დასახელება, გამოცემის ადგილი, წელი, ჟურნალის №, პირველი და ბოლო გვერდები). მონოგრაფიის შემთხვევაში მიუთითეთ გამოცემის წელი, ადგილი და გვერდების საერთო რაოდენობა. ტექსტში კვადრატულ ფხიხლებში უნდა მიუთითოთ ავტორის შესაბამისი N ლიტერატურის სიის მიხედვით. მიზანშეწონილია, რომ ციტირებული წყაროების უმეტესი ნაწილი იყოს 5-6 წლის სიღრმის.

9. სტატიას თან უნდა ახლდეს: ა) დაწესებულების ან სამეცნიერო ხელმძღვანელის წარდგინება, დამოწმებული ხელმოწერითა და ბეჭდით; ბ) დარგის სპეციალისტის დამოწმებული რეცენზია, რომელშიც მითითებული იქნება საკითხის აქტუალობა, მასალის საკმაობა, მეთოდის სანდოობა, შედეგების სამეცნიერო-პრაქტიკული მნიშვნელობა.

10. სტატიის ბოლოს საჭიროა ყველა ავტორის ხელმოწერა, რომელთა რაოდენობა არ უნდა აღემატებოდეს 5-ს.

11. რედაქცია იტოვებს უფლებას შეასწოროს სტატია. ტექსტზე მუშაობა და შეჯერება ხდება საავტორო ორიგინალის მიხედვით.

12. დაუშვებელია რედაქციაში ისეთი სტატიის წარდგენა, რომელიც დასაბეჭდად წარდგენილი იყო სხვა რედაქციაში ან გამოქვეყნებული იყო სხვა გამოცემებში.

აღნიშნული წესების დარღვევის შემთხვევაში სტატიები არ განიხილება.

Содержание:

Rahardjo H.E., Ückert S., Maerker V., Bannowsky A., Kuczyk M.A., Kedia G.T. STIMULATION OF THE CYCLIC AMP/GMP SIGNALLING ENHANCES THE RELAXATION OF ISOLATED HUMAN DETRUSOR SMOOTH MUSCLE ACHIEVED BY PHOSPHODIESTERASE INHIBITORS	7
Styopushkin S., Chaikovskiy V., Chernylovskiy V., Sokolenko R., Bondarenko D. POSTOPERATIVE HEMORRHAGE AS A COMPLICATION OF A PARTIAL NEPHRECTOMY: FREQUENCY, FEATURES AND MANAGEMENT.....	12
Бурьянов А.А., Лыходий В.В., Задниченко М.А., Соболевский Ю.Л., Пшеничный Т.Е. КЛИНИЧЕСКАЯ ОЦЕНКА РЕЗУЛЬТАТОВ ХИРУРГИЧЕСКОГО ЛЕЧЕНИЯ ПАЦИЕНТОВ С ДЕГЕНЕРАТИВНЫМИ ПОВРЕЖДЕНИЯМИ КОРНЯ МЕДИАЛЬНОГО МЕНИСКА	20
Чернооков А.И., Рамишвили В.Ш., Долгов С.И., Николаев А.М., Атаян А.А., Белых Е.Н. СОВРЕМЕННАЯ СТРАТЕГИЯ ЛЕЧЕНИЯ БОЛЬНЫХ С РЕЦИДИВАМИ ВАРИКОЗНОЙ БОЛЕЗНИ ПОСЛЕ ЭНДОВАЗАЛЬНЫХ ВМЕШАТЕЛЬСТВ.....	26
Babaskin D., Litvinova T., Babaskina L., Krylova O., Savinova O., Winter E. EFFECT OF ELECTRO- AND ULTRAPHONOPHORESIS OF THE PHYTOCOMPLEX ON MICROCIRCULATORY AND BIOCHEMICAL PARAMETERS IN PATIENTS WITH KNEE JOINT OSTEOARTHRITIS	34
Japaridze Sh., Lomidze L., Nakhutsrishvili I., Davituliani V., Kekelidze I. APPLICATION OF ANTIBIOTIC-CONTAINING EAR DROPS IN TREATMENT OF ACUTE OTITIS MEDIA.....	41
Sevbitov A., Emelina E., Khvatov I., Emelina G., Timoshin A., Yablokova N. EFFECT OF SMOKING STEAM COCKTAILS ON THE HARD TISSUES OF THE ORAL CAVITY	44
Borysenko A., Dudnikova M. CLINICAL RATIONALE OF CHOOSING A TOOTH-BLEACHING AGENT	48
Kladnichkin I., Ivanov S., Bekreev V., Salata A., Trufanov V. METHODOLOGY FOR CONSISTENT COPYING OF THE OVERDENTURE RESTORATION PARAMETERS FOR DENTAL IMPLANT PROSTHESIS IN THE TREATMENT OF TOTAL EDENTIA.....	51
Гоциридзе К.Э., Кинтрая Н.П., Гогия Т.Э., Надареишвили Л.Н. ИММУННЫЕ НАРУШЕНИЯ И ИХ РОЛЬ В ПРЕРЫВАНИИ БЕРЕМЕННОСТИ.....	57
Sirko A., Mizyakina K., Chekha K. POST-TRAUMATIC HEADACHE. CURRENT VIEWS ON PATHOPHYSIOLOGICAL MECHANISMS OF DEVELOPMENT AND CLINICAL SPECIFICS (REVIEW)	60
Fedorenko S., Onopriienko I., Vitomskiy V., Vitomska M., Kovelska A. INFLUENCE OF A PSYCHOTYPE OF A PATIENT WITH MUSCULOSKELETAL DISORDER ON THE DEGREE OF WORK DISABILITY.....	66
Krylov A., Khorobrykh T., Petrovskaya A., Khmyrova S., Agadzhyanov V., Khusainova N. ROLE OF THROMBODYNAMICS GLOBAL COAGULATION TEST IN IMPROVING TREATMENT RESULTS IN PATIENTS WITH CORONAVIRUS INFECTION AT A COVID-19 HOSPITAL	72
Petrov V., Molozhavenko E., Ivashina E., Sozonov A., Baksheev E. LASER THERMAL ABLATION OF BENIGN THYROID NODULES AS AN EFFECTIVE, SAFE AND MINIMALLY INVASIVE METHOD FOR TREATING NODULAR GOITER (REVIEW)	79
Gavrtsyuk V., Merenkova I., Vlasova N., Vychenko O. CLINICAL FACTORS ASSOCIATED WITH THE RISK OF PULMONARY SARCOIDOSIS RELAPSE	84
Дорош Д.Н., Лядова Т.И., Волобуева О.В., Попов Н.Н., Сорокина О.Г., Огнивенко Е.В. КЛИНИКО-ИММУНОЛОГИЧЕСКИЕ ОСОБЕННОСТИ ГЕРПЕСВИРУСНЫХ ЗАБОЛЕВАНИЙ НА ФОНЕ ВИЧ.....	89

Ivakhniuk T., Ivakhniuk Yu. INTESTINAL MICROBIOTA IN ALZHEIMER'S DISEASE	94
Lazashvili T., Silagadze T., Kapetivadze V., Tabukashvili R., Maglapheridze Z., Kuparadze M. ACTION OF SIMVASTATIN IN IMPROVING COGNITIVE FUNCTIONS IN VASCULAR DEMENTIA.....	98
Kolinko L., Shlykova O., Izmailova O., Vesnina L., Kaidashev I. SIRT1 CONTRIBUTES TO POLARIZATION OF PERIPHERAL BLOOD MONOCYTES BY INCREASING STAT6 EXPRESSION IN YOUNG PEOPLE WITH OVERWEIGHT AND LOW-RISK OBESITY	102
Акимов М.А., Политова А.С., Пекарский С.П., Коваленко В.В., Телефонко Б.М. ПСИХИЧЕСКОЕ РАССТРОЙСТВО КАК ОБЯЗАТЕЛЬНЫЙ МЕДИЦИНСКИЙ КРИТЕРИЙ ОГРАНИЧЕННОЙ ВМЕНЯЕМОСТИ	113
Жармаханова Г.М., Сырлыбаева Л.М., Кононец В.И., Нурбаулина Э.Б., Байкадамова Л.И. МОЛЕКУЛЯРНО-ГЕНЕТИЧЕСКИЕ АСПЕКТЫ РАЗВИТИЯ МЕТИЛМАЛОНОВОЙ АЦИДУРИИ (ОБЗОР)	118
Zhvania M., Kvezereli-Kopadze M., Kutubidze T., Kapanadze N., Gordeladze M., Iakobashvili A., Nakhutsrishvili E. COVID-19 AND CHILDREN: COMPLICATIONS AND LATE OUTCOMES.....	124
Tukhtiyeva N., Dossanov B., Sakalouski A., Syzdykbayev M., Zhunussov Y. METHODS OF TREATMENT OF LEGG - CALVÉ - PERTHES DISEASE (REVIEW)	127
Shengelia M., Burjanadze G., Koshoridze M., Kuchukashvili Z., Koshoridze N. STRESS-AFFECTED Akt/mTOR PATHWAY UPREGULATED BY LONG-TERM CREATINE INTRAPERITONEAL ADMINISTRATION.....	134
Morar I., Ivashchuk A., Bodaya V., Domanchuk T., Antoniv A. FEATURES OF GRANULATION TISSUE MORPHOLOGY AROUND THE NET ALLOTRANSPLANT WHEN APPLYING POSTOPERATIVE RADIATION THERAPY	139
Харисова Н.М., Смирнова Л.М., Кузьмин А.Ф., Рыспаева Г.К., Лепесбаева Г.А. ОСОБЕННОСТИ РАЗВИТИЯ РЕПРОДУКТИВНОЙ СИСТЕМЫ ПРИ ИСПОЛЬЗОВАНИИ ГЕНЕТИЧЕСКИ МОДИФИЦИРОВАННЫХ ИСТОЧНИКОВ (ЭКСПЕРИМЕНТАЛЬНОЕ ИССЛЕДОВАНИЕ).....	146
Nikolaishvili M., Nanobashvili Z., Mitagvaria N. RADON HORMESIS IN EPILEPTIC PATHOGENESIS AND PREDICTORS OF OXIDATIVE STRESS.....	152
Ходели Н.Г., Чхаидзе З.А., Шенгелия О.С., Сонгулашвили Д.П., Инаури Н.А. СОВЕРШЕНСТВОВАНИЕ ПЕРФУЗИОННОГО ПОТОКА НАСОСОВ КРОВИ.....	158
Гнатюк М.С., Татарчук Л.В., Крицак М.Ю., Коноваленко С.О., Слабый О.Б., Монастырская Н.Я. МОРФОМЕТРИЧЕСКАЯ ОЦЕНКА ОСОБЕННОСТЕЙ РЕМОДЕЛИРОВАНИЯ КРОВЕНОСНЫХ СОСУДОВ СЕМЕННИКОВ ПРИ АРТЕРИАЛЬНОЙ ГИПЕРТЕНЗИИ В МАЛОМ КРУГЕ КРОВООБРАЩЕНИЯ В ЭКСПЕРИМЕНТЕ	163
Goncharuk O., Savosko S., Petriv T., Medvediev V., Tymbaliuk V. QUANTITATIVE HISTOLOGICAL ASSESSMENT OF SKELETAL MUSCLE HYPOTROPHY AFTER NEUROTOMY AND SCIATIC NERVE REPAIR IN RATS	169
Sharashenidze T., Shvelidze Kh., Tsimakuridze M., Turabelidze-Robaqidze S., Buleishvili M., Sanikidze T. ROLE OF β -ADRENOCEPTORS IN REGULATION OF ERYTHROCYTES' RHEOLOGICAL FUNCTIONS (REVIEW)	173
Afanasieva M., Stoianov M., Kuli-Ivanchenko K., Ivanchenko A., Shotova-Nikolenko A. VACCINATION: STATE-IMPLEMENTED MEDICO-SOCIAL AND LEGAL MEASURES.....	176
Булеца С.Б., Заборовский В.В., Менджул М.В., Пирого И.С., Тымчак В.В., Стойка А.В. ПРАВОВАЯ ЗАЩИТА И ОСОБЕННОСТИ ПРИМЕНЕНИЯ ТЕХНОЛОГИЙ ВИРТУАЛЬНОЙ РЕАЛЬНОСТИ В МЕДИЦИНЕ	180
Осмолян В.А., Домбровская Е.Н., Хорошенко О.В. УЧАСТИЕ ВРАЧА В ДОПРОСЕ НЕСОВЕРШЕННОЛЕТНЕГО ЛИЦА КАК ОБЯЗАТЕЛЬНАЯ ПРАВОВАЯ НОРМА В ЗАКОНОДАТЕЛЬСТВЕ	186

нейными фибриллами, свидетельствующий об иммунных реакциях при Covid-инфекции, чем и следует объяснить положительный эффект кортикостероидной терапии в лечении этих больных. 22 (36%) пациента из 60 госпитализированы, остальные наблюдались амбулаторно.

На основе вышеизложенного следует заключить, что даже при асимптомном течении Covid-инфекции у детей могут наблюдаться осложнения и формирование т.н. синдрома позднего Covid-a, что диктует необходимость тщательного обследования этих больных и наблюдения в динамике.

რეზიუმე

Covid-19 და ბავშვები: გართულებები და შორეული გამოსავალი

მ.ჟვანია, მ.კვეხერელი-კოპაძე, თ.კუტუბიძე, ნ.კაპანაძე, მ.გორდელაძე, ა.იაკობაშვილი, ე.ნახუცრიშვილი

თბილისის სახელმწიფო სამედიცინო უნივერსიტეტი, გ. ჟვანიას სახ. პედიატრიის აკადემიური კლინიკა, საქართველო

2019 წლის დეკემბრიდან Covid-19 მსოფლიოს მედიკოსების, მათ შორის პედიატრების, გამოწვევა გახდა. ინფიცირებულ ბავშვთა უმეტესობაში დაავადება მსუბუქი ფორმით ვლინდება, ან ხასიათდება სუბკლინიკური მიმდინარეობით. ზოგიერთ შემთხვევაში შეიძლება განვითარდეს დაავადების მოგვიანებითი მიმე კლინიკური სურათი, მულტისისტემური სინდრომის და სხვა გართულებების სახით.

თბილისის სახელმწიფო სამედიცინო უნივერსიტეტის გ. ჟვანიას სახ. აკადემიურ პედიატრიულ კლინიკაში 2020-2021 წლებში დაკვირვების ქვეშ იმყოფებოდა 60 ბავშვი, რომელთაც აღენიშნებოდათ

პოსტკოვიდური გართულებები და ე.წ. გვიანი Covid-19 სინდრომი. 60 ინფიცირებული ბავშვიდან 51 (85%) დაავადებამდე იყო ჯანმრთელი. 32 (53,3%) იყო 5 წლამდე ასაკის. ინფიცირებულ ბავშვებში დომინირებდა მამრობითი სქესი - 33 (55%). 60 პაციენტს 1,5-2 თვით ადრე მომართვამდე გადატანის დადებითი ტიტრით დადასტურდა. კლინიკურად მათ აღენიშნათ ვასკულოპათია, იმუნური თრომბოციტოპენია, რკინადეფიციტური ანემია, კოაგულოპათია, პნევმონია-ატელექტაზი, პირველადი დიაბეტი, ძირითადი დაავადების გამწვავება - ართრალგია, ართრიტი და ზოგადი სიმპტომატიკა ძილის დარღვევის, სისუსტისა და თავბრუსხვევის სახით. ცალკე უნდა აღინიშნოს მულტისისტემური ანთებითი სინდრომი 8 (13%) ბავშვში, კავასაკის დაავადების კლინიკური ნიშნებით (კან-ლორწოვან-ლიმფური კვანძების სინდრომი), ჰექტიური ცხელებით, პოლისეროზიტით, ჰეპატოსპლენომეგალიით, ანთების მარკერების მაღალი მაჩვენებლებით, ტენდენციით ჰიპერკოაგულაციისაკენ. ერთ პაციენტს აღენიშნა კორონარული არტერიის ანევრიზმა. 3 შემთხვევაში ANA ტიტრი მომატებული იყო 1:640-მდე, ასევე დაფიქსირდა ბირთვების ციტოპლაზმური ფლუორესცენცია ხაზოვანი ფიბრილებით, რაც Covid-ინფექციის იმუნურ რეაქციაზე მიანიშნებს და კორტიკოსტეროიდული თერაპიის ეფექტურობას ამართლებს. 60-დან 22 (36%) პაციენტი იყო ჰოსპიტალიზებული, დანარჩენი 38 იმყოფებოდა ამბულატორული მეთვალყურეობის ქვეშ.

ყოველივე ზემოთქმულიდან გამომდინარე ავტორებს გამოტანილი აქვთ დასკვნა, რომ ბავშვებში Covid-19 ინფექციის ასიმპტომური მიმდინარეობის დროსაც კი შესაძლებელია აღინიშნოს გართულებები და ე.წ. გვიანი კოვიდის სინდრომის ჩამოყალიბება, რაც ამ პაციენტების სრულყოფილ გამოკვლევასა და დინამიკაში დაკვირვების აუცილებლობას მოითხოვს.

METHODS OF TREATMENT OF LEGG - CALVÉ - PERTHES DISEASE (REVIEW)

¹Tuktiyeva N., ²Dossanov B., ³Sakalouski A., ¹Syzdykbayev M., ¹Zhunussov Y.

¹NCJSC "Semey Medical University", Department of children's surgery and orthopedics;

²NCJSC "Astana Medical University", Department of children's surgery, Nur-Sultan, Republic of Kazakhstan;

³Republican Scientific and Practical Center of Traumatology and Orthopedics, Minsk, Belarus

More than a century has passed and the problem of treating Legg-Calvé-Perthes Disease (LCPD) remains relevant. Children between the ages of 2 and 12 tend to get sick. Formation and progression causes a violation of blood supply to the femoral head over time. Subsequently, this leads to changes in the femoral head, metaphysis, epiphyseal plate, and coxal cavity [32]. The flaccidity of the femur epiphysis is deformed during load and leads to deformation of the femoral head due to uneven load transmission [35].

Elimination of pain and symptoms, restoration of the range of movement of the hip, and the retention of the femoral head in the coxal cavity are the targets of the treatment [55].

Dissatisfaction with the results of treatment has led to the existence of many conservative and prompt treatments [12,18,23,26]

Purpose of review - to review the various existing methods of conservative and operative therapies of LCPD, which include the elimination of pain and symptoms, the restoration of the range of hip movements, and the retention of the femoral head in the coxal cavity.

Material and methods. Literature search was conducted through PubMed and Google Scholar using keywords Legg - Calvé - Perthes Disease, Perthes disease; operative therapy, conservative therapy, childhood hip disorder. 5921 publications were initially identified. Articles were published between December 1971 and August 2020.

Result and discussion. Treatment of Legg-Calve-Perthes disease is not lost in relevance in time and presents serious problems, due to long rehabilitation and a large percentage of unsatisfactory outcomes, despite the existence of various successful treatments. That promotes the search for more rational approaches to the therapy and perfect surgical interventions. There are two great areas for LCPD therapy: conservative and surgical.

1. Conservative therapy. The conservative therapy is divided into three main areas:

1. Unloading the joint and immobilization of the limb for a very long time.
2. Orthoses and application of a plaster bandage to immerse the femoral head into the coxal cavity.

Improving blood supply or increasing the strength of the femoral head - physiotherapy, balneotherapy, hyperbaric oxygenation, the use of glycosaminoglycans and bisphosphonates, plasma therapy, platelets enriched, epidural analgesia, complex conservative therapy, etc [48].

1.1. Terms of conservative therapy

According to Wiig et al. non-surgical treatment is provided to children under the age of 6 or in case of lesion of the A lateral pillar [71].

The determining points for conservative therapy are the age of the patient and the stage of the disease - the group I and partly the II according to the classifications Catterall, Salter & Thompson [17].

The state of full "immersion" of the femoral head into the coxal cavity stops the progression of deformation of the head and prevents the development of subluxation, as well as helps to correct the deformation [15]. In order to unload the joint for a long time, until reaching maturity of bone tissue, used - orthopedic unloading corsets, plaster bandages, stretching cuff or adhesive and other methods [7,10,67].

In herring studies and co-authors in 2004, they examined the results of treatment of about 400 patients over the age of 6 with LCPD, using 5 types of therapy. According to the results, there is no statistically significant difference between therapeutic gymnastics and observation, plaster bandage or braces. The unsatisfactory outcome reached 20 %, according to the Stulberg classification. According to P.G. Petrie, I. Bitenca & B. Curtis, they have demonstrated excellent results using of plaster bandages and orthoses, the negative result was only in 9-17 % of cases [11].

Different methods of conservative therapy are effective in the early stages, with a slight lesion of the segment. In case of extensive damage to the femoral head, as well as in the contracture of a hip joint, it is necessary to use another treatment [4].

1.1. Methods based on the principles of improving blood supply or increasing the strength of the femoral head

1.0.1. Bisphosphonates

Bisphosphonates are known to be inhibitors of osteoclast resorption of the bone tissue. There are data from experimental studies, which say that bisphosphonates reduce bone resorption and deformation of the head. But the distribution of bisphosphates depends on the blood supply and their influence is limited accordingly [37]. In an experimental study on animals, the anatomical structure of the femoral head was observed and the expressed decrease of osteoclasts [36]. Biophosphates are achieved not only inhibiting the process of osteoclast resorption, but also accelerating osteoblast activity [35].

1.0.2. Acupuncture

There is a description of clinical case T. Set, in the absence of the effect of brace treatment in a 12-year-old boy, 196 sessions

of classical and laser acupuncture were received over a period of 4 years. After 2 years of acupuncture according to X-ray data, the recovery was 90%. After 6 years, there was a complete restoration of the anatomical structure of the femoral head [62].

1.0.3. Antioxidants

Lobashov V.V. cites an approximate successful complex treatment with antioxidants (mexidol). During the studies, 26 children had received antioxidant therapy (intramuscular, ointment form, intraosseous injections). A positive result was achieved in 84.6 % of cases and unsatisfactory 15.4% [48].

1.0.4. Epidural analgesia

Prolonged epidural analgesia in Legg-Calvé-Perthes Disease improves the local tissue trophics, helping to eliminate spasms in ischemia, reducing pain syndrome. There was also a slowdown in the flow of aseptic necrosis. The method allowed to delay total hip replacement for up to 6 years-old-age [2,3].

1.0.5. Physiotherapy

In the complex treatment of the Legg-Calvé-Perthes disease, intralesional electrophoresis of proteolytic enzymes was proposed. The author believed that these manipulations reduce increased intraosseous pressure, help to germinate the additional feeding vessels, and together with grafts stimulate bone regeneration in the head of the hip [56].

In studies comparing different treatments, physiotherapy has been applied for children with mild disease. There were the following patient characteristics: children with hip necrosis were less than 50% (Catterall Group 1 or 2). As well as children under six years old with necrosis of the femoral head were more than 50%, who have a good coating of the femoral head (>80%) [20].

For patients with a light course, physiotherapy can improve joint range, muscle strength and joint dysfunction. The physiotherapy treatment included: 1) passive mobilization for stretching the muscles of the affected hip; 2) exercises with lifting straight legs to strengthen the muscles of the hip, participating in bending, flexing, lifting, and engaging the thigh muscles. They started with isometric exercises, and after eight sessions - isotonic exercises. Balance training is first on a stable terrain and then on unstable.

There are scientifically sound recommendations for postoperative LCPD treatment in children between the ages of 3 and 12 years and evidence-based recommendations for conservative LCPD treatment in children between the ages of 3 and 12. These studies are largely based on the "local consensus" of LCPD team members from Cincinnati Children's Hospital Medical Center. These recommendations provide evidence on the methods of physiotherapy, postoperative and conservative management. Recommendations are offered: exercises to improve balance and gait and intervention to reduce pain.

Children with LCPD should avoid hip overload. Gait training to unload the hip can be an integral component of conservative treatment for children with LCPD.

Non-surgical corset treatment is a reliable alternative to surgical treatment of LCPD between the ages of 6 and 8. However, they could not have known whether the good results were due to the corset or the result of a good prognosis for these patients [20,58].

1. Operative therapy

The history of operative therapy for ischemic necrosis of the femur head has passed several stages. In the beginning, palliative methods of operative therapy were used to stimulate reparative processes in the epiphys area of the femur. The method of tunneling, drilling of the femur head and its neck by Beck has become widespread [31].

Assuming in the future possible early development of osteoarthritis, many orthopedists recognize the benefits of more active, operative tactics to treat this disease [11,40,54,68,70]

In case of violation of anatomical ratios at the stage of exodus, reconstructive operations are carried out.

Operative therapy of Legg-Calvé-Perthes Disease is carried out with an unfavorable prognosis in order to improve the volume of movements in the joint, restore limb length, cupping pain syndrome [16, 70].

The long-term results of operating interventions are more than modest. In the evaluation of 59 patients operated between 1959 and 1974, in 2002 36 % of patients had osteoarthritis, and in 2010 – 50 % of patients only [61].

In 2012, Nhu-An has compared the results of conservative and operative therapies of patients. The meta-analysis showed that operative therapy was more likely to contribute to the formation of a spherical congruent femur head than conservative treatment in patients of 6-years-old age and older. The patients younger than 6 years-old age the results of operative and conservative therapy were identical [55]. Positive results to restore the anatomical spherical shape of the femoral head are dependent on age: the younger the child at the time of the disease, the more effective the therapy is [69].

In Kim HK research, Su PH. the outcomes of Legg-Calvé-Perthes Disease were studied in 66 patients with an average follow-up period of 20. 4 years (16. 3 to 24. 5 years). The most common complaints in patients were pain syndrome, developing arthrosis and ongoing hip dysfunction. Total arthroplasty was performed by 3 patients and 1 patient carried out pelvic osteotomy. 24 patients suffered from severe osteoarthritis. Only 14 patients had no signs of osteoarthritis [38].

Endoprosthesis replacement at an early age in patients with LCPD contributes to lower survival rates, according to the study [34].

A number of authors propose a combination of operative and conservative therapies for LCPD [50].

2.1. Femoral or pelvis osteotomy

Femoral or pelvis osteotomy is indicated in children over 8 years of age. Femoral varus osteotomy and anonymous Salter-type osteotomy give good results. However, in severe forms, it is preferable to combine these two methods or perform a triple pelvic osteotomy. Surgery is now much less common than before, as it is only effective in patients with the disease of lateral support of group B or B/C with the onset after the age of eight. In other situations, therapeutic abstinence is recommended [47].

The main task of osteotomy is to ensure the centering of the femoral head and congruence of joint surfaces, recently osteotomy has become one of the common operative methods. The optimal time to perform a femoral dethroxy-varisizing osteotomy is considered to be the stage of avascular necrosis or the initial period of the fragmentation stage [14, 29]. Many authors advocate inter-reversive correctional femoral osteotomy [19, 41, 44, 53, 57].

In the stage of the exodus of Legg-Calvé-Perthes Disease with an expressed shortening of the limb and hypertrophy of the greater trochanter developed femoral rotation-longing interverent osteotomy [16, 15, 28]. Salter-type surgery has led to the possibility of improving the congruence of joint surfaces with existing changes in angular parameters, as well as a significant improvement in the blood supply to the joint compared to other types of osteotomy. The pelvic component was corrected with various types of pelvic osteotomy in combination or without interference with the proximal part of the femur [59, 60, 58]

There are surgical interventions in the form of acetabuloplasty, supplemented, if applied, by interverent osteotomy. In most cases, orthopedists prefer to use Salter-type pelvic osteotomy and modifications of triple pelvic osteotomy such as Steel, Tonnis, Chiari, rotational acetabular osteotomy, Ganz, Bernese. This group of surgical interventions has a number of advantages: the necrosis center avoids load, there is reliable fixation of bone fragments, restoration of biomechanical beneficial relationships of joint surfaces, reduction of the period of postoperative rehabilitation [13,47].

Kyung Soon Park et al. speaks of the successful use of modified Salter-type osteotomy, produced in 30 cases with satisfactory long-term clinical and radiological re-examinations [42].

Asep Santoso et al. gives examples of the results of successful treatment of osteotomy dometatasis of the pelvis combined with trochanteric advancement for sequelae of Perthes' disease. Good results of treatment of acetabular dysplasia due to Perthes disease were obtained in the middle or long-term period. Pain relief was received in 13 out of 14 (92.8%) patients in the post-operative period. From good to excellent functionality, the result was obtained from 10 out of 14 (71.4%) patients. The average score on the Harris scale improved from 63 to 84 ($p < 0.05$) in the last phase of the observation. Improvements in lame gait were observed in 10 out of 14 (71.4%) patients [5].

Yi cites the results of the use of Bernese-type triple pelvic osteotomy with a modified Smith-Peterson anterior approach, which combines periacetabular and triple osteotomy. The author believes that this technique is an alternative method of treatment in the elderly people. Not only does it provide a large correction of the coxal cavity, but it also provides good biomechanical stability [75].

Negative outcome with the development of deforming osteoarthritis, with the right selected and carried out operational treatment, reaches 20%. That requires a subsequent complete joint replacement [2]. Surgical treatment in the form of femoral osteotomy is a factor complicating subsequent hip replacement. It is known from the statistics that bilateral form occurs in 10-20 % of cases, which also makes it difficult to carry out operative therapy while maintaining the patient's mobility [44]. On the other hand, the difficulty of choosing a conservative treatment method causes a positive outcome of the disease in 30-50% of cases, with the restoration of the spherical head of the femur, without significant statistical differences of the chosen method [27].

J. Lehman et al. speak about the successful use of arthroplasty in the treatment of Legga-Calvé-Perthes disease [46]. Luo and co-author cite the assessment of 10-year results in uncemented monoblock total arthroplasty of the hip 71 patients (88 hip joints) were examined. The data suggest that the trunk of the monoblock can lead to satisfactory results - the restoration of clinical function, the improvement of X-ray evaluation, the restoration of normal limb length, reduction of complications and increased survival among patients [76].

As the healing potential decreases with age, patients with LCPD should receive treatment corresponding to their age group. Conservative treatment is usually applied to patients under 6.0 years-old of age from the beginning, and surgical treatment is recommended for those over 8.0 years of age, but it remains unclear what is best for patients between the ages of 6.0 and 8.0 years. The purpose of this retrospective study was to compare the results of Salter-type osteotomy and conservative therapy in this age group. As a result, there was no significant difference in hip pain and joint mobility between groups in this age group [33].

The long-term results of The Chiari-type osteotomy in Legg-Calve-Perthes disease in children with Catterall III or IV type have also been studied. Chiari-type osteotomy with 15 degrees for Legg-Calve-Perthes disease in children with Catterall III or IV type can effectively lower the index of the coxal cavity and can help to change the shape of the femur and then further improve the clinical effects [77].

2.2. Valgus or regimented osteotomy

Improves the spinal mechanism of a teenager's hip with the effects of Perthes' disease. Proper surgery, when the hip is still in the pre-arytritic stage, restores function and protects the young hip from early degenerative changes.

Eid cites the results of treatment of 12 adolescents with a combined intra-articular/injected impinge due to the effects of Perthes disease in the form of coxamagna was carried out osteochondroplasty at the junction of the femoral head and cervix and the relative lengthening of the femoral cervix by the spiritual transfer of a large twithlet, coxaplana, coxabrevis, with the predominantly large vernea. In all cases, safe surgical access to a dislocated hip was performed. Removal of the narrow arc of pathological load due to the impingement is the main advantage of the proposed surgical technique. Other benefits include increased leverage and the restoration of hip motion range with normalization of load conditions, and therefore the future development of degenerative arthritis and the expected need for future joint replacement surgery can be prevented or delayed. Vascularization of the femur head is well supported thanks to the proven safety of the surgical access presented [24].

Osteostimulative interventions in the form of multiple tunnelization of the femoral head and cervix with the introduction of allografts were performed in patients with X-ray indicators of the hip joint close to the age norm. Over time, the operative intervention was improved in the form of transplantation on the nourishing vascular-muscular leg in the upper acetabular area [22].

There was a low efficiency of tunnelization of the head of the femur with the introduction of surgery transplants, in view of the absence of trophies of free bone grafts vascularization was ensured and often observed progression of the disease [63].

Decompressive operations are becoming popular, providing a hip discharge, reducing intraarticular pressure, weakening muscle contractures and reducing the load on the affected femoral head. To this end the following were applied: the tenotomy of the iliac-lum, leading and subspinal muscles, cross-cutting in several places of the wide fascia of the hip followed by adhesive or skeletal extremity of the limb. Since 1993, external fixation devices such as G.A. have been used to unload the hip joint Ilizarov [65]. In 1965 Axer A. proposed varizing detorsion osteotomy as an alternative method in the treatment of Legg-Calve-Perthes disease [8].

In order to unload the most damaged quadrant of the head, a method of inter-flexion-warising femoral osteotomy is developed and introduced into clinical practice, in which, unlike deprofessionalizing, another element is performed - the flexion of the proximal part of the femur. In the development of the surgical intervention Bernbeck R's operation was taken as the basis, the main principle of which is to change the angle of the slope of the fragments of the obliquely sawed cylinder when they rotate on the axis, perpendicular plane of the section [52]. On the basis of this principle, a method of intervening oblique osteotomy of the femur has been developed, allowing to achieve a multi-axis simultaneous correction of the proximal end of the femur [39].

Leclerc at al. it was concluded that the best effect was achieved by putting the lower limb 30 degrees in the outside and limiting

the stretch time to no more than two weeks, otherwise prolonged stretching contributes to a negative effect [45].

Long-term observations showed that varus osteotomy of the femur was an effective treatment for patients with Perthes disease between the ages of 6 and 8 with Herring B and C hip lesion at the onset of the disease. Hip congruence was obtained by femoral varus osteotomy, and closed wedge osteotomy gave more favorable results to CE angle [6].

Barakat et al. offers a closed repository in older children an effective and reproducible method of treatment in the hands of an experienced pediatric orthopedic surgeon, subject to careful observation and careful study of possible complications and their treatment, including the possibility of a timely transition to an open reposition [9].

For children over 6 years-old age diagnosed with femoral head necrosis, more than 50% proximal varus osteotomy has produced significantly better results than orthoses and physiotherapy [72].

2.3. Hip arthroscopy

New treatments for mechanical symptoms and/or femoral acetabular impingement have emerged. Arthroscopic treatment of the effects of LCPD eased pain symptoms and improved the range of motion, making arthroscopic treatment a good option for treating the effects of LCPD [73].

Kanatli in his study demonstrates improvements in functional outcomes and quality of life for patients who have undergone hip arthroscopy due to LCPD. It is believed that hip arthroscopy, a minimally invasive procedure, may play an important role in the LCPD treatment algorithm, especially in patients with severe pain and mechanical symptoms [66].

Hip arthroscopy was performed to 19 children with Legg-Calve-Perthes disease in the Shigeo Suzuki et al. As a result, the growth of the synovial shell was expressed both in the swivel pit and over the inner wall of the capsule. Hypervascularization of the twirling lip was observed at all stages of the disease. Microscopic hyperplasia of synovial lining cells was observed, but inflammatory changes in synovial tissue were invisible in the early stages of the disease. Although endothelial vascular cell hypertrophy was observed at the late stage of the disease, it was not distinct at the initial stage or stage of fragmentation. Joint pain decreased after irrigation during arthroscopy [64].

Freeman et al. presented the results of the arthroscopy of Legg-Calve-Perthes disease and reflected that it played an important role in the treatment of painful consequences. It is often possible to expect successful results with minimal morbidity. Reducing symptoms and improving quality of life are reasonable expectations, although these data do not suggest that hip arthroscopy alters the natural history of the disease [25].

Hip arthroscopy is a safe procedure even for patients with an immature skeleton. It turns out to be effective for improving functional disorders caused between the deformed head of the femur and the coxal cavity, or some intraarticular focal problems in hip diseases in children and adolescents. However, this does not mean that hip arthroscopy can prevent or delay the development of osteoarthritis [21].

2.4. Arthrodiastase of the hip joint

What can be done with regard to indications of the use of arthrodiastase in the active stage with the late onset of Perthes disease or as a rescue procedure in later cases with debilitating deformity. There is insufficient evidence to suggest whether it is superior to other interventions and may be combined with hip or pelvis osteotomy as stage 2 surgery. More randomized control is needed to be tested to compare with other treatments [74].

Carlos Augusto Malheiros Luzo and co-authors presented preliminary results of treatment of patients with Legga-Calve-Perthes disease using hip monolateral external fixer applied to the hip for 18 patients. Among the patients were 13 men and 5 women, the average age is 8.5, from 5 to 13 years-old. All patients had one-sided hip damage: nine on the right and nine on the left. The results were evaluated in adulthood using clinical and radiological criteria. Results: all patients improved joint mobility, pain relief was achieved in 88.9% of cases. The reossification of the epiphys of the femur occurred during the first three months of treatment. The hips operated on at the stage of necrosis of the disease did not pass the stage of fragmentation, which led to a reduction in the period of the disease. The results were 77.8% satisfactory and 22.2% unsatisfactory improved joint mobility. The use of arthrodiastase technique at the stage of necrosis or at the stage of fragmentation (active phase of the disease) gave satisfactory results from treatment. Arthrodiastase of the hip joint with the use of a one-sided external retainer in the active stages of the disease improves joint mobility. The use of arthrodiastase technique at the stages of necrosis and fragmentation (active stages of the disease) gives satisfactory results in the treatment of the disease [49]. Maxwell and co-author investigated the effect of arthrodiastase on the preservation of the head of the femur in older children with Perthes disease. Arthrodiastase was used for about four months. The initial criterion for evaluation was the degree of epiphysal collapse at the end of the fragmentation phase. One of the 15 treated hips and nine of the 30 control hips showed a loss of height of 50% or more lateral epiphysal column on the front of the X-ray (Herring classification). From Lauenstein's point of view, one of the treated thighs and 19 control thighs showed a loss of height of at least 50 % of the anterior epiphysal column. Complications of arthrodiastase included a pin infection in most hip joints, temporary joint stiffness in two and a broken finger on two. The final result will be known when all patients and the control group reach the maturity of the skeleton [51].

Arthrodiastase is a minimally invasive technique useful in the treatment of late-onset Perthes disease. When applied early, it improves clinical outcomes and retains the shape of the head. However, it is not free of complications that can occur in 8 out of 10 patients treated with this technique [1].

Arthrodiastase, hinge or fixed using an external ring retainer in combination with the lead muscle tenotomy, is an excellent and reliable treatment for Perthes disease with late onset, when the prognosis is usually poor and traditional treatments are unreliable [30].

Arthrodiastase of the hip joint with the release of soft tissues can make a good contribution to the treatment of Legg-Calve-Perthes disease. This method of treatment has many advantages, such as simplicity of technique, minimal frequency of complications, short period of hospitalization, correction of shortening because it increases the length of the limb, and a higher level of acceptable results than would be expected compared to other methods. Reduces upper and lateral sub-raving and provides better X-ray spherical sphericity of the head of the femur. In addition, it does not distort the anatomy of the pelvis or the proximal part of the femur; it can also be used in older children who usually have poor prognosis. Distraction treatment is not limited to hip stiffness, femur or sub-infection, and can be used when other treatments are not suitable [43].

Conclusion. When choosing a treatment method for Legg-Calve-Perthes disease, it is necessary to focus on the specific clinical case. As well as take into account the age of the patient

at the time of the beginning of the pathological process, the stage of the disease, the size and location of the lesion in the head of the femur, predictive anatomical and functional factors.

REFERENCES

1. Aguado-Maestro I., Abril J. C., Bañuelos D. A., García A. M. Hip arthrodiastasis in Legg-Calvé-Perthes disease. *Rev Esp Cir Ortop Traumatol*, 2016, 60(4), 243-50.
2. Akhtyamov I.F. & Lobashov V.V. Aseptic necrosis of the femoral head in children, the options of conservative treatment. *Bulletin of modern clinical medicine*, 2014, 2, 40-44.
3. Akhtyamov I.F., Anisimov O.G. & Kovalenko A.N. New way to treat early forms of avascular necrosis of the femoral head (preliminary report). *Bulletin of Traumatology and Orthopedics m. N.N. Priorova*, 2011, 1, 33-37.
4. Akhtyamov I.F., Lobashov V.V., Anisimov O.G. A method of treating the early stages of Perthes' disease (preliminary Report). *Journal of Traumatology and Orthopedics of Russia* 2014; 3: 122-128.
5. Asep S., Pramod S. I., Ik-Sun Ch., Young-Rok Sh., Kyung-Soon P. & Taek-Rim Y. Treatment outcome of dome osteotomy of the pelvis combined with trochanteric advancement for sequelae of Perthes' disease. *Acta Orthopaedica et Traumatologica Turcica*, 2018, 52, 216-221.
6. Atilla C. Long-term follow-up results of femoral varus osteotomy in the treatment of Perthes disease, and comparison of open-wedge and closed-wedge osteotomy techniques A retrospective observational study. *Medicine (Baltimore)*. 2020, 99 (7): e19041. PMID: 32049801doi: 10.1097/MD.00000000000019041
7. Aurelio G., Martinier M. D., Stuart L. et al. The Weight-Bearing Abduction Brace for the Treatment of Perthes Disease. *J. Bone Jt. Surg*, 1992, 74(1), 12-21.
8. Axer A. Subtrochanteric osteotomy in the treatment of Perthes' disease: a preliminary report. *J Bone Joint Surg. Br*, 1965, 47, 489-499.
9. Barakat A. S., Zein A. B. & Arafa A. S., et al. Closed reduction with or without adductor tenotomy for developmental dysplasia of the hip presenting at walking age. *Curr Orthop Pract*, 2017, 28 (2), 195-199. doi 10.1097/BCO.0000000000000478.
10. Barsukov, D. B. Perthes disease. *Terra medica nova* 2009; 3: 24-30.
11. Beaulé P., E. & Antoniadis. Patient selection and surgical technique for surface arthroplasty of the hip. *Orthop.Clin. North Am*, 2005, 36 (2), 177-185.
12. Benjamin J. Management of Perthes' disease. *Indian J Orthop*, 2015, 49(1), 10-16.
13. Bhuyan B. K. Early outcomes of one-stage combined osteotomy in Legg-Calve-Perthes diseases, *Indian J Orthop*, 2016, 50 (2), 183-194.
14. Cartier P., Dagher F. & Morel L. La osteotomie de varisation dans la maladie de Legg-Calve-Perthes. *Rev. Chir. Orthop.* 1976, 62, 27-41.
15. Catterall A. Legg-Calve-Perthes Syndrome. *Clin.Orthop*, 1981, 158, 41-52.
16. Catterall A. Natural history, classification and X-ray signs in Legg-Calve-Perthes disease. *Acta orthop. Helv*, 1980, 46 (4), 346-351.
17. Catterall A. The natural history of Perthes disease. *J. Bone Joint Surg. Br*, 1971, 53, 37-52.
18. Catterall A., Pringle J. & Byers P., D., et al. Perthes disease: is the epiphysal infarction complete? *J Bone Joint Surg. Br*, 1982, 64(3), 276-281.

19. Cosimo G., Frizziero P., & Turra S. Prognostic value of Catterall and Herring classification in Legg-Calve-Perthes disease: follow-up to skeletal maturity of 32 patients. *J. Pediatr. Orthop.* 2002, 22 (3), 345-349.
20. Cincinnati Children's Hospital Medical Center. Evidence-based clinical care guideline for Conservative Management of Legg-Calve-Perthes Disease. Guideline, 2011, 39. <https://www.cincinnatichildrens.org/-/media/cincinnati%20childrens/home/service/j/anderson-center/evidence-based-care/recommendations/type/legg-calve-perthes%20disease%20guideline%2039>
21. Chaemoon L., Tae-Joon Cho., Chang H. Sh., In H. Ch., Won J., Y. Functional Outcomes of Hip Arthroscopy for Pediatric and Adolescent Hip Disorders. *Clin Orthop Surg*, 2020, 12 (1), 94-99. <https://doi.org/10.4055/cios.2020.12.1.94>
22. Griffin P.P. Legg-Calve-Perthes disease: treatment and prognosis. *Optop. Clin. North Am*, 1999, 11 (1), 127-139.
23. Gu Y. Can an enlarged acetabulum cover the femoral head Well in Legg-Calve-Perthes disease? *J. Pediatr Orthop*, 1999, 8 (3), 173-176.
24. Eid M. A. Hip preservation surgery for adolescents and young adults with Post-Perthes Sequelae. *Acta Orthop Belg*. 2016, 82 (4), 821-828. PMID: 29182124.
25. Freeman C., R., Jones, K., Byrd, J., W. Hip arthroscopy for Legg-Calve-Perthes disease: minimum 2-year follow-up. *Arthroscopy*, 2013, 29, 666-674.
26. Guarniero R. Results of femoral varus osteotomy in the treatment of Legg-Calve-Perthes disease. *Rev. Hosp.Clin. Fas. Med Sao Paulo*. 1997, 52 (3), 132-135.
27. Herring J., Kim H. & Browne R. Legg — Calvé — Perthes disease: Part II. Prospective multicenter study of the effect of treatment on outcome. *J. Bone Joint Surg. Am*, 2004, 86 (10), 2121-34.
28. Herring J. A., Neustadt J. B., Williams J., J. Early J. S. & Browne R.H. The lateral pillar classification of Legg-Calve-Perthes disease. *J Pediatr.Orthop*, 1992, 12 (2), 143-150.
29. Imatani J. Coxa magna after open reduction for developmental dislocation of the hip. *J. Pediatr. Orthop*, 1995, 15 (3), 337-341.
30. José B., V. Comparison between innominate osteotomy and arthrodistraction as a primary treatment for Legg-Calvé-Perthes disease: a prospective controlled trial. *IntOrthop*. 2012, 36 (9), 1899–1905. doi: 10.1007/s00264-012-1598-2PMCID: PMC3427447 PMID: 22810494
31. Joseph D. L., Perry L. S., Pascual-Garrido C., Jeffery J. N. & John C. C. Open Reduction and Internal Fixation for the Treatment of Symptomatic Osteochondritis Dissecans of the Femoral Head in Patients With Sequelae of Legg-Calve-Perthes Disease. *Journal of Pediatric Orthopaedics*, 2018, 40(3), 120-128. doi: 10.1097/BPO.0000000000001192
32. Joseph B. Natural history of early onset and late onset Legg-Calve-Perthes disease. *J. Pediatr Orthop*, 2011, 31, 152–155.
33. Kaneko H., Kitoh H., Mishima K., Matsushita M., Hattori T., Noritake K., Ishiguro N. & Yoshihashi Y. Comparison of surgical and nonsurgical containment methods for patients with Legg-Calvé-Perthes disease of the onset ages between 6.0 and 8.0 years: Salter osteotomy versus a non-weight-bearing hip flexion-abduction brace. *J Pediatr Orthop B*. 14. *J Pediatr Orthop B*, 2020, 29(6), 542-549. Doi: 10.1097/BPB.0000000000000710. <https://www.ncbi.nlm.nih.gov/pubmed/31856043>
34. Karim Z., M., John J. C. & José A. M. Primary total hip arthroplasty for Legg-Calve-Perthes syndrome: 20 year follow-up study. *The Iowa Orthopedic Journal*, 2018, 38, 197-202.
35. Kim HK. W. Pathophysiology and new strategies for the treatment of Legg-Calve-Perthes disease. *J Bone Joint Surg. Am*, 2012, 94, 659-669.
36. Kim HK., Morgan-Bagley S. & Kostenuik P. RANKL inhibitor: a novel strategy to decrease femoral head deformity after ischemic osteonecrosis. *J Bone Joint Mineral Res*, 2006, 21, 1246-1254.
37. Kim HK. W., Randall T. S. & Bian H. et al. Ibandronate for prevention of femoral head deformity after ischemic necrosis of the capital femoral epiphysis in immature pigs. *J Bone Joint Surg. Am*, 2005, 3, 550-557.
38. Kim HK. & Su PH. Development flattening and apparent fragmentation following ischemic necrosis of the capital femoral epiphysis in a piglet model. *J Bone Joint Surg. Am*, 2009, 91, 2903-3014.
39. Kitakoji T. Femoral varus osteotomy in Legg-Calve-Perthes disease: points of operation to prevent residual problems. *J. Pediatr. Orthop*, 1999, 19 (1), 76-81.
40. Koob T. J., Pringle D. & Gedbaw E. et al. Biomechanical properties of bone and cartilage in growing femoral head following ischemic osteonecrosis. *J Orthop Res*, 2007, 25, 750-757.
41. Kwang W. P., Chastity A. R., Rey An N. G., Tae Wan K. & Hae Ryong Song. Extent of physeal involvement in Leg-Calve-Perthes disease. *Inter.Orthop*. 2014, 38 (11), 2303-2308.
42. Kyung S. P., Kyu J., Hong Y. Y., Kamolhuja E. & Taek R. Y. Long-term Results of Modified Salter Innominate Osteotomy for Legg-Calve-Perthes Disease. *Clinics in Ortopedic Surgery*, 2017, 9, 397-404. <https://doi.org/10.4055/cios.2017.9.4.397>
43. Lakloul M. A., Hosny & Gamal A. Hinged distraction of the hip joint in the treatment of Perthes disease. *Journal of Pediatric Orthopaedics B*, 2012, 21(5), 386–393. doi:10.1097/bpb.0b013e328354b0ab
44. Larson A. N., Sucato D. J. & Herring J. A., et al. A Prospective Multicenter Study of Legg-Calve-Perthes Disease. Functional and Radiographic Outcomes of Nonoperative Treatment at a mean follow-up of twenty years. *J Bone Joint Surg. Am*. 2012, 94, 584-592.
45. Leclerc J., Laville J. M. & Salmeron F. Review of orthopedic surgery and repair of the motor system, 2006, 92 (8), 741-746.
46. Lehmann T. G., Engesaeter I., Laborie L. B, Lie S. A, Rosendahl K. & Engesaeter L. B. Total hip arthroplasty in young adults, with focus on Perthes' disease and slipped capital femoral epiphysis: Followup of 540 subjects reported to the Norwegian Arthroplasty Register during 1987-2007. *Acta Orthop*, 2012, 83, 159–64.
47. Leroux J., Abu A. S. & Lechevallier J. Legg-Calve-Perthes disease. *Indian J. Orthop*. 2018, 104, 107-112. Doi: 10.4103/0019-5413.177581.
48. Lobashov V.V. Conservative treatment of disease Legg - Calvet – Perthes. *Modern art of medicine*. 2013, 5 (13), 14-19.
49. Luzo C. A. M., Guarniero R., Montenegro N. B., Godoy Junior R. M. Initial experience of use of an articulated external fixator in treating Legg-Calvé-Perthes disease by means of arthrodiastasis during the active phase of the disease. *Revista Brasileira de Ortopedia (English Edition)*, 2016, 51(3), 337–345. doi:10.1016/j.rboe.2016.04.005
50. Mallet C., Abitan A., Vidal C., Holvoet L., Mazda K. & Simon B. I. Management of osteonecrosis of the femoral head in children with sickle cell disease: results of conservative and operative treatments at skeletal maturity. *J Child Orthop*, 2018, 12, 47-54.
51. Maxwell S. L., Lappin K. J., Kealey W. D., McDowell B. C. & Cosgrove A. P. Arthrodiastasis in Perthes' disease - Preliminary results. *The Bone & Joint Journal*, 2004, 86 (2), 244-60. DOI: 10.1302/0301.620X.86B2.14284
52. Menshikov T. I. & Maltseva L.V. Features of ultrasound diagnosis of the initial manifestations of Legg-Calvet-Perthes disease in children. *International Journal of Experimental Education*, 2015, 11 (1), 18-23.

53. Milani C., Dobashi E., T. Arthrogram in Legg-Calve-Perthes disease. *J Pediatr.Orthop*, 2011, 31, 156-162.
54. Miyamoto Y., Matsuda T., Kitoh H., et al. A recurrent mutation in type II collagen gene causes Legg-Calve-Perthes disease in a Japanese family. *Hum Genet*. 2007, 121, 625-629.
55. Nhu-An T. N., Guy K. G., Jessica McCourt B. & Charles T. M. Operative versus nonoperative treatments for Legg-Calvé-Perthes disease: a meta-analysis. *J PediatrOrthop*, 2012, 32(7), 697-705. doi: 10.1097/BPO.0b013e318269c55d.
56. Ogarev E. V. Formation of the proximal femur in children and adolescents. *Vestn. traumatology and orthopedics them. N.N. Priorov*, 2006, 1, 51-56.
57. Perry D. C., Machin D. M., Pope D. & Bruce C. E. et al. Racial and geographic factors in the incidence of Legg-Calve-Perthes disease: a systematic review. *Am J Epidemiol*, 2012, 175 (3), 159-166.
58. Pinto H. A. & Peterson T. et al. Magnetic resonance imaging in earle diagnosis of Legg-Calve-Perthes disease. *J. Pediatr. Orthop*, 1989, 9, 19-22.
59. Ponseti I. V. Legg-Calve-Perthes disease; observations on pathological changes into two cases. *J Bone Joint Surg. Am*, 1956, 38, 739-750.
60. Pringle D., Koob T. J. & Kim H. K. Indentation properties of growing femoral head following ischemic necrosis. *J Orthop, Pes*, 2004, 22, 122-130.
61. Salter R. B. & Thompson G. H. Legg-Calve-Perthes Disease: the prognostic significance of the subchondral fracture and a two-group classification of the femoral head involvement. *J. Bone Joint Surg. Am*, 1984, 66 (4), 479-489.
62. Set T. Management of Legg — Calve — Perthes disease with acupuncture: a case report. *Acupunct. Med*, 2013, 1, 105-112.
63. Shapovalov V. M. & Tikhilov R. M. Surgical treatment of degenerative-dystrophic diseases of the hip joint. *Vestnik Surgery them I.I. Grekova*, 1999, 6, 44-47.
64. Shigeo S., Yoshitaka K., Yoichi S., Toru F., Kengo F. & Yoshihito N. Arthroscopy in 19 children with perthes' disease: Pathologic changes of the synovium and the joint surface P. *Acta Orthopaedica Scandinavica*, 1994, 69, 581-584. <https://doi.org/10.3109/17453679408994608>
65. Teplen'kii M.P., Parfenov E.M. Surgical treatment of children with severe forms of Perthes disease. *Geniy ortopedii*, 2013, 1, 32-35.
66. Ulunay K., Tacettin A., Mustafa O., Muhamme, B., A. & Mehmet C. Hip arthroscopy for Legg-Calvé-Perthes disease in paediatric population. *ActaOrthopaedica et Traumatologica Turcica*, 2019, 53 (3), 203-208.
67. Veselovsky Yu. A. Pathogenesis and early conservative treatment of the initial stages of osteochondropathy of the femoral head in children. *Orthoped, trauma*, 1989, 4, 4-7.
68. Weinstein S. L. Developmental Hip Dysplasia and Dislocation. *Pediatric Orthopedics*, 1996, 2, 903-950.
69. Weinstein S. L. Legg-Calvé-Perthes syndrome. In: Lowell & Winter' *Pediatric Orthopaedy*. Philadelphia, etc: Lippincott-Raven Pub, 1996, 2 (4), 951-991.
70. Westhoff B., Petersomann A., Hirsch M. A., Willers R. & Krauspe R. Computerized gait analysis in Legg-Calve-Perthes disease - analysis of the frontal plane. *Gait and Posture*, 2006, 24 (1), 196-202.
71. Wiig O., Svenningsen S., Terjesen T. Legg-Calvé-Perthes disease. *Tidsskr. Nor. Laegeforen*, 2011, 20, 131 (9-10), 946-953.
72. Wiig O., Terjesen T., Svenningsen S. Prognostic factors and outcome of treatment in Perthes' disease. *J Bone Joint Surg*, 2008, 90-B (10), 1364-1371.

73. Woo-Yong L., Deuk-Soo H., Pil-Sung K. & Lon Zh. Outcomes in patients with late sequelae (healed stage) of Legg-Calvé-Perthes disease undergoing arthroscopic treatment: retrospective case series. *Hip Int*. 2018, 28 (3), 302-308. doi: 10.5301/hipint.5000563
74. Yasser H., I., Mohamed A, A., Kersh L., Hesham F. Arthrodiastasis in the management of Perthes disease: a systematic review. *Journal of Pediatric Orthopaedics B*, 2019, Vol XXX, 1-6
75. Yi Q. L., Hong W. Xu., Theddy S., Qing H. Z., Yuanzhong L., Wei D. Ch., Jing Ch. L. & Federic C. Bernese type triple pelvic osteotomy through a single incision in children over five years: a retrospective study of twenty eight cases. *International Orthopaedics* 2018, 42, 2961-2968. <https://doi.org/10.1007/s00264-018-3946-3>
76. Ze-Yu L., Hao-Yang W., Duan W., Hui P., Fu-Xing P., & Zong-Ke Zhou. Musculoskeletal Disorders. *BMC*, 2017, 18, 386. DOI 10.11. 1186/s12891-017-1748-1
77. Zang J. T., Wang Y. H., Feng W., Liu J. G., Li D. S., Zhao X. Y., Gao H., Li C. S. Long-term outcomes of Chiari osteotomy for Legg-Calvé-Perthes disease in children. *Zhongguo Gu Shang*. 2019, 32(8), 696-700. doi: 10.3969/j.issn.1003-0034.2019.08.004.

SUMMARY

METHODS OF TREATMENT OF LEGG-CALVÉ-PERTHES DISEASE (REVIEW)

¹Tuktıyeva N., ²Dossanov B., ³Sakalouski A.,
¹Syzdykbayev M., ¹Zhunussov Y.

¹NCJSC "Semey Medical University", Department of children's surgery and orthopedics; ²NCJSC "Astana Medical University", Department of children's surgery, Nur-Sultan, Republic of Kazakhstan; ³Republican Scientific and Practical Center of Traumatology and Orthopedics, Minsk, Belarus

This Literature Review presents various treatments, including operative and conservative therapies, of Legg - Calvé - Perthes Disease. The problem is relevant because of the prevalence of the disease. The authors presented a review of literature, which managed to classify the main methods of treatment by the principles of action, practical application, and presented the interpretation of the effectiveness of modern research from the point of view of evidence-based medicine.

Keywords: Legg-Calvé-Perthes disease; Perthes disease; operative therapy; conservative therapy; childhood hip disorder.

РЕЗЮМЕ

МЕТОДЫ ЛЕЧЕНИЯ БОЛЕЗНИ ЛЕГГА-КАЛЬВЕ-ПЕРТЕСА (ОБЗОР)

¹Туктиева Н.А., ²Досанов Б.А., ³Соколовский О.А.,
¹Сыздыкбаев М.К., ¹Жунусов Е.Т.

¹НАО «Медицинский университет Семей», кафедра детской хирургии и ортопедии; ²НАО «Медицинский университет Астана», кафедра детской хирургии Нур-Султан, Республика Казахстан; ³Республиканский научно-практический центр травматологии и ортопедии, Минск, Беларусь

В литературном обзоре проанализированы различные методы лечения (оперативные, консервативные) болезни Легга —

Кальве — Пертеса. Проблема актуальна ввиду распространенности заболевания. На основании анализа и синтеза текущей и ретроспективной научной литературы авторами рассмотрены

и классифицированы основные методы лечения по принципам действия, практическому применению, а также дана оценка эффективности с точки зрения доказательной медицины.

რეზიუმე

ლევა-კალვე-პერტესის დაავადების მკურნალობის მეთოდები (მიმოხილვა)

¹ნ.ტუკტიევა, ²ბ.დოსანოვი, ³ო.სოკოლოვსკი, ⁴მ.სიხდიკბაევი, ⁵ე.ჟუნუსოვი

¹სემეის სამედიცინო უნივერსიტეტი, ბავშვთა ქირურგიისა და ორთოპედიის კათედრა;
²ასტანის სამედიცინო უნივერსიტეტი, პედიატრიული ქირურგიის კათედრა, ნურ-სულთან,
ყაზახეთის რესპუბლიკა; ³ტრავმატოლოგიისა და ორთოპედიის რესპუბლიკური
სამეცნიერო-პრაქტიკული ცენტრი, მინსკი, ბელორუსია

ლიტერატურის მიმოხილვაში გაანალიზებულია თანამედროვე და რეტროსპექტიული სამეცნიერო ლიტერატურა ლევა-კალვე-პერტესის დაავადების სამკურნალო მეთოდების შესახებ. განხილულია მკურნალობის ოპერაციული და კონსერვატი-

ული მეთოდები, რომლებიც კლასიფიცირებულია მკურნალობის, მოქმედების პრინციპების და პრაქტიკული გამოყენების გათვალისწინებით. განხილულია მასალის ანალიზის და სინთეზის შედეგად ავტორებს მოცემული აქვთ მეთოდების ეფექტურობის შეფასება.

STRESS-AFFECTED Akt/mTOR PATHWAY UPREGULATED BY LONG-TERM CREATINE INTRAPERITONEAL ADMINISTRATION

Shengelia M., Burjanadze G., Koshoridze M., Kuchukashvili Z., Koshoridze N.

Ivane Javakhishvili Tbilisi State University, Department of Biology, Faculty of Exact and Natural Sciences, Georgia

Oxidative stress is known to be characterized by significant alterations in metabolic processes, namely changes in the hormonal status, decreased energy metabolism and antioxidant status, as well as quantitative changes in enzyme activity and signalling molecules, which, in turn, affect transcription and translation processes [20,24]. Several compounds can prevent these processes. Among them is the Creatine (Cr; α -N-methylguanidinoacetic acid), which can be found in almost all mammals. It is primarily concentrated in muscle and brain. It participates in the Cr/CK/PCr system, is actively involved in energy metabolism, and its deficiency is associated with a decline in many physical and cognitive functions [2,9,13]. It is believed that the primary mechanism of action of Creatine (Cr) is its participation in the energy storage processes. Besides, various experiments also confirmed its neuromodulatory and neuroprotective properties [4,17]. Cr synthesized in nerve cells functions as a signalling molecule. In particular, it can activate some signalling pathways and, in this way, regulate energy metabolism, influencing growth, proliferation and viability of the cell [1,14]. In the brain, Creatine is most concentrated in the regions associated with learning processes and memory (such as Hippocampus, Pyramidal neurons of the cortex, Purkinje cells of the cerebellum). It is assumed that these areas are also marked with high ATP metabolism [21].

Cr is not only established to be synthesized by neurons, but it is also suggested to be delivered peripherally through the blood-brain barrier [2,4]. Exogenous Cr showed its neuroprotective properties in the number of neurological diseases such as Parkinson's disease, Huntington's disease, Amyotrophic lateral

sclerosis (ALS), head injuries [3]. Quantitative changes in Cr have also been shown in various psychiatric disorders, such as depression [1,5].

Recent data have further revealed the antioxidant properties of Cr [14,29]. Observations have shown that lipid peroxidation processes are down-regulated as antioxidant enzymes are activated in muscle and central nervous system (CNS), during Cr supplementation [14]. Such alterations might be caused by several stressors, such as long-term violation of natural circadian rhythm [26]. This kind of stress is usually accompanied by a change in antioxidant and energy metabolism - resulting in ATP deficiency, the brain's energy potential and functional deterioration, cell viability reduction, stimulation of pro-apoptotic processes, and variation in ion content [32].

Considering the above mentioned, the purpose of our investigation was to study complications in energy metabolism in the hippocampus under stress caused by long-term disturbance of circadian rhythm and the preventive action of Cr administered exogenously.

Material and method. Experiments were conducted on 200–250 gr male Wistar rats. The animals were divided into three main groups before the experiment:

- (1) G1 – control group – was kept in a common cage under natural conditions (dark/light ratio = 10/14);
- (2) G2 – stressed group – individuals were maintained in individual cages in the darkness (dark/light ratio = 23.5/0.5) for 30 days;
- (3) G3 – Cr-treated stressed - individuals were maintained in individual cages in the dark (dark/light ratio = 23.5/0.5) for 30