

# GEORGIAN MEDICAL NEWS

---

ISSN 1512-0112

NO 10 (355) Октябрь 2024

---

ТБИЛИСИ - NEW YORK



ЕЖЕМЕСЯЧНЫЙ НАУЧНЫЙ ЖУРНАЛ

Медицинские новости Грузии  
საქართველოს სამედიცინო სიახლენი

## GEORGIAN MEDICAL NEWS

Monthly Georgia-US joint scientific journal published both in electronic and paper formats of the Agency of Medical Information of the Georgian Association of Business Press.  
Published since 1994. Distributed in NIS, EU and USA.

**GMN: Georgian Medical News** is peer-reviewed, published monthly journal committed to promoting the science and art of medicine and the betterment of public health, published by the GMN Editorial Board since 1994. GMN carries original scientific articles on medicine, biology and pharmacy, which are of experimental, theoretical and practical character; publishes original research, reviews, commentaries, editorials, essays, medical news, and correspondence in English and Russian.

GMN is indexed in MEDLINE, SCOPUS, PubMed and VINITI Russian Academy of Sciences. The full text content is available through EBSCO databases.

**GMN: Медицинские новости Грузии** - ежемесячный рецензируемый научный журнал, издаётся Редакционной коллегией с 1994 года на русском и английском языках в целях поддержки медицинской науки и улучшения здравоохранения. В журнале публикуются оригинальные научные статьи в области медицины, биологии и фармации, статьи обзорного характера, научные сообщения, новости медицины и здравоохранения. Журнал индексируется в MEDLINE, отражён в базе данных SCOPUS, PubMed и ВИНТИ РАН. Полнотекстовые статьи журнала доступны через БД EBSCO.

**GMN: Georgian Medical News** – საქართველოს სამედიცინო სიახლენი – არის ყოველთვიური სამეცნიერო სამედიცინო რეცენზირებადი ჟურნალი, გამოიცემა 1994 წლიდან, წარმოადგენს სარედაქციო კოლეგიისა და აშშ-ის მეცნიერების, განათლების, ინდუსტრიის, ხელოვნებისა და ბუნებისმეტყველების საერთაშორისო აკადემიის ერთობლივ გამოცემას. GMN-ში რუსულ და ინგლისურ ენებზე ქვეყნდება ექსპერიმენტული, თეორიული და პრაქტიკული ხასიათის ორიგინალური სამეცნიერო სტატიები მედიცინის, ბიოლოგიისა და ფარმაციის სფეროში, მიმოხილვითი ხასიათის სტატიები.

ჟურნალი ინდექსირებულია MEDLINE-ის საერთაშორისო სისტემაში, ასახულია SCOPUS-ის, PubMed-ის და ВИНТИ РАН-ის მონაცემთა ბაზებში. სტატიების სრული ტექსტი ხელმისაწვდომია EBSCO-ს მონაცემთა ბაზებიდან.

### WEBSITE

[www.geomednews.com](http://www.geomednews.com)

## К СВЕДЕНИЮ АВТОРОВ!

При направлении статьи в редакцию необходимо соблюдать следующие правила:

1. Статья должна быть представлена в двух экземплярах, на русском или английском языках, напечатанная через **полтора интервала на одной стороне стандартного листа с шириной левого поля в три сантиметра**. Используемый компьютерный шрифт для текста на русском и английском языках - **Times New Roman (Кириллица)**, для текста на грузинском языке следует использовать **AcadNusx**. Размер шрифта - **12**. К рукописи, напечатанной на компьютере, должен быть приложен CD со статьей.

2. Размер статьи должен быть не менее десяти и не более двадцати страниц машинописи, включая указатель литературы и резюме на английском, русском и грузинском языках.

3. В статье должны быть освещены актуальность данного материала, методы и результаты исследования и их обсуждение.

При представлении в печать научных экспериментальных работ авторы должны указывать вид и количество экспериментальных животных, применявшиеся методы обезболивания и усыпления (в ходе острых опытов).

4. К статье должны быть приложены краткое (на полстраницы) резюме на английском, русском и грузинском языках (включающее следующие разделы: цель исследования, материал и методы, результаты и заключение) и список ключевых слов (key words).

5. Таблицы необходимо представлять в печатной форме. Фотокопии не принимаются. **Все цифровые, итоговые и процентные данные в таблицах должны соответствовать таковым в тексте статьи**. Таблицы и графики должны быть озаглавлены.

6. Фотографии должны быть контрастными, фотокопии с рентгенограмм - в позитивном изображении. Рисунки, чертежи и диаграммы следует озаглавить, пронумеровать и вставить в соответствующее место текста **в tiff формате**.

В подписях к микрофотографиям следует указывать степень увеличения через окуляр или объектив и метод окраски или импрегнации срезов.

7. Фамилии отечественных авторов приводятся в оригинальной транскрипции.

8. При оформлении и направлении статей в журнал МНГ просим авторов соблюдать правила, изложенные в «Единых требованиях к рукописям, представляемым в биомедицинские журналы», принятых Международным комитетом редакторов медицинских журналов - <http://www.spinesurgery.ru/files/publish.pdf> и [http://www.nlm.nih.gov/bsd/uniform\\_requirements.html](http://www.nlm.nih.gov/bsd/uniform_requirements.html) В конце каждой оригинальной статьи приводится библиографический список. В список литературы включаются все материалы, на которые имеются ссылки в тексте. Список составляется в алфавитном порядке и нумеруется. Литературный источник приводится на языке оригинала. В списке литературы сначала приводятся работы, написанные знаками грузинского алфавита, затем кириллицей и латиницей. Ссылки на цитируемые работы в тексте статьи даются в квадратных скобках в виде номера, соответствующего номеру данной работы в списке литературы. Большинство цитированных источников должны быть за последние 5-7 лет.

9. Для получения права на публикацию статья должна иметь от руководителя работы или учреждения визу и сопроводительное отношение, написанные или напечатанные на бланке и заверенные подписью и печатью.

10. В конце статьи должны быть подписи всех авторов, полностью приведены их фамилии, имена и отчества, указаны служебный и домашний номера телефонов и адреса или иные координаты. Количество авторов (соавторов) не должно превышать пяти человек.

11. Редакция оставляет за собой право сокращать и исправлять статьи. Корректур авторам не высылаются, вся работа и сверка проводится по авторскому оригиналу.

12. Недопустимо направление в редакцию работ, представленных к печати в иных издательствах или опубликованных в других изданиях.

**При нарушении указанных правил статьи не рассматриваются.**

## REQUIREMENTS

Please note, materials submitted to the Editorial Office Staff are supposed to meet the following requirements:

1. Articles must be provided with a double copy, in English or Russian languages and typed or computer-printed on a single side of standard typing paper, with the left margin of 3 centimeters width, and 1.5 spacing between the lines, typeface - **Times New Roman (Cyrillic)**, print size - 12 (referring to Georgian and Russian materials). With computer-printed texts please enclose a CD carrying the same file titled with Latin symbols.

2. Size of the article, including index and resume in English, Russian and Georgian languages must be at least 10 pages and not exceed the limit of 20 pages of typed or computer-printed text.

3. Submitted material must include a coverage of a topical subject, research methods, results, and review.

Authors of the scientific-research works must indicate the number of experimental biological species drawn in, list the employed methods of anesthetization and soporific means used during acute tests.

4. Articles must have a short (half page) abstract in English, Russian and Georgian (including the following sections: aim of study, material and methods, results and conclusions) and a list of key words.

5. Tables must be presented in an original typed or computer-printed form, instead of a photocopied version. **Numbers, totals, percentile data on the tables must coincide with those in the texts of the articles.** Tables and graphs must be headed.

6. Photographs are required to be contrasted and must be submitted with doubles. Please number each photograph with a pencil on its back, indicate author's name, title of the article (short version), and mark out its top and bottom parts. Drawings must be accurate, drafts and diagrams drawn in Indian ink (or black ink). Photocopies of the X-ray photographs must be presented in a positive image in **tiff format**.

Accurately numbered subtitles for each illustration must be listed on a separate sheet of paper. In the subtitles for the microphotographs please indicate the ocular and objective lens magnification power, method of coloring or impregnation of the microscopic sections (preparations).

7. Please indicate last names, first and middle initials of the native authors, present names and initials of the foreign authors in the transcription of the original language, enclose in parenthesis corresponding number under which the author is listed in the reference materials.

8. Please follow guidance offered to authors by The International Committee of Medical Journal Editors guidance in its Uniform Requirements for Manuscripts Submitted to Biomedical Journals publication available online at: [http://www.nlm.nih.gov/bsd/uniform\\_requirements.html](http://www.nlm.nih.gov/bsd/uniform_requirements.html)  
[http://www.icmje.org/urm\\_full.pdf](http://www.icmje.org/urm_full.pdf)

In GMN style for each work cited in the text, a bibliographic reference is given, and this is located at the end of the article under the title "References". All references cited in the text must be listed. The list of references should be arranged alphabetically and then numbered. References are numbered in the text [numbers in square brackets] and in the reference list and numbers are repeated throughout the text as needed. The bibliographic description is given in the language of publication (citations in Georgian script are followed by Cyrillic and Latin).

9. To obtain the rights of publication articles must be accompanied by a visa from the project instructor or the establishment, where the work has been performed, and a reference letter, both written or typed on a special signed form, certified by a stamp or a seal.

10. Articles must be signed by all of the authors at the end, and they must be provided with a list of full names, office and home phone numbers and addresses or other non-office locations where the authors could be reached. The number of the authors (co-authors) must not exceed the limit of 5 people.

11. Editorial Staff reserves the rights to cut down in size and correct the articles. Proof-sheets are not sent out to the authors. The entire editorial and collation work is performed according to the author's original text.

12. Sending in the works that have already been assigned to the press by other Editorial Staffs or have been printed by other publishers is not permissible.

**Articles that Fail to Meet the Aforementioned  
Requirements are not Assigned to be Reviewed.**

## ავტორთა საქურაღებოლ!

რედაქციაში სტატიის წარმოდგენისას საჭიროა დაიცვათ შემდეგი წესები:

1. სტატია უნდა წარმოადგინოთ 2 ცალად, რუსულ ან ინგლისურ ენებზე დაბეჭდილი სტანდარტული ფურცლის 1 გვერდზე, 3 სმ სიგანის მარცხენა ველისა და სტრიქონებს შორის 1,5 ინტერვალის დაცვით. გამოყენებული კომპიუტერული შრიფტი რუსულ და ინგლისურენოვან ტექსტებში - **Times New Roman (Кириллица)**, ხოლო ქართულენოვან ტექსტში საჭიროა გამოვიყენოთ **AcadNusx**. შრიფტის ზომა – 12. სტატიას თან უნდა ახლდეს CD სტატიით.

2. სტატიის მოცულობა არ უნდა შეადგენდეს 10 გვერდზე ნაკლებს და 20 გვერდზე მეტს ლიტერატურის სიის და რეზიუმეების (ინგლისურ, რუსულ და ქართულ ენებზე) ჩათვლით.

3. სტატიაში საჭიროა გაშუქდეს: საკითხის აქტუალობა; კვლევის მიზანი; საკვლევი მასალა და გამოყენებული მეთოდები; მიღებული შედეგები და მათი განსჯა. ექსპერიმენტული ხასიათის სტატიების წარმოდგენისას ავტორებმა უნდა მიუთითონ საექსპერიმენტო ცხოველების სახეობა და რაოდენობა; გაუტკივარებისა და დაძინების მეთოდები (მწვავე ცდების პირობებში).

4. სტატიას თან უნდა ახლდეს რეზიუმე ინგლისურ, რუსულ და ქართულ ენებზე არანაკლებ ნახევარი გვერდის მოცულობისა (სათაურის, ავტორების, დაწესებულების მითითებით და უნდა შეიცავდეს შემდეგ განყოფილებებს: მიზანი, მასალა და მეთოდები, შედეგები და დასკვნები; ტექსტუალური ნაწილი არ უნდა იყოს 15 სტრიქონზე ნაკლები) და საკვანძო სიტყვების ჩამონათვალი (key words).

5. ცხრილები საჭიროა წარმოადგინოთ ნაბეჭდი სახით. ყველა ციფრული, შემაჯამებელი და პროცენტული მონაცემები უნდა შეესაბამებოდეს ტექსტში მოყვანილს.

6. ფოტოსურათები უნდა იყოს კონტრასტული; სურათები, ნახაზები, დიაგრამები - დასათაურებული, დანომრილი და სათანადო ადგილას ჩასმული. რენტგენოგრაფიების ფოტოასლები წარმოადგინეთ პოზიტიური გამოსახულებით **tiff** ფორმატში. მიკროფოტოსურათების წარწერებში საჭიროა მიუთითოთ ოკულარის ან ობიექტივის საშუალებით გადიდების ხარისხი, ანათალებების შედეგების ან იმპრეგნაციის მეთოდი და აღნიშნოთ სურათის ზედა და ქვედა ნაწილები.

7. სამამულო ავტორების გვარები სტატიაში აღინიშნება ინიციალების თანდართვით, უცხოურისა – უცხოური ტრანსკრიპციით.

8. სტატიას თან უნდა ახლდეს ავტორის მიერ გამოყენებული სამამულო და უცხოური შრომების ბიბლიოგრაფიული სია (ბოლო 5-8 წლის სიღრმით). ანბანური წყობით წარმოდგენილ ბიბლიოგრაფიულ სიაში მიუთითეთ ჯერ სამამულო, შემდეგ უცხოელი ავტორები (გვარი, ინიციალები, სტატიის სათაური, ჟურნალის დასახელება, გამოცემის ადგილი, წელი, ჟურნალის №, პირველი და ბოლო გვერდები). მონოგრაფიის შემთხვევაში მიუთითეთ გამოცემის წელი, ადგილი და გვერდების საერთო რაოდენობა. ტექსტში კვადრატულ ფხიხლებში უნდა მიუთითოთ ავტორის შესაბამისი N ლიტერატურის სიის მიხედვით. მიზანშეწონილია, რომ ციტირებული წყაროების უმეტესი ნაწილი იყოს 5-6 წლის სიღრმის.

9. სტატიას თან უნდა ახლდეს: ა) დაწესებულების ან სამეცნიერო ხელმძღვანელის წარდგინება, დამოწმებული ხელმოწერითა და ბეჭდით; ბ) დარგის სპეციალისტის დამოწმებული რეცენზია, რომელშიც მითითებული იქნება საკითხის აქტუალობა, მასალის საკმაობა, მეთოდის სანდოობა, შედეგების სამეცნიერო-პრაქტიკული მნიშვნელობა.

10. სტატიის ბოლოს საჭიროა ყველა ავტორის ხელმოწერა, რომელთა რაოდენობა არ უნდა აღემატებოდეს 5-ს.

11. რედაქცია იტოვებს უფლებას შეასწოროს სტატია. ტექსტზე მუშაობა და შეჯერება ხდება საავტორო ორიგინალის მიხედვით.

12. დაუშვებელია რედაქციაში ისეთი სტატიის წარდგენა, რომელიც დასაბეჭდად წარდგენილი იყო სხვა რედაქციაში ან გამოქვეყნებული იყო სხვა გამოცემებში.

აღნიშნული წესების დარღვევის შემთხვევაში სტატიები არ განიხილება.

Nino Chichua, Giuli Margishvili, Grigol Dzodzuashvili, Rusudan Ivanishvili, Vladimer Margvelashvili. EVALUATING ORAL AND MAXILLOFACIAL HEALTH CHALLENGES IN INTRAVENOUS DRUG USERS: A CROSS-SECTIONAL STUDY OF DRUG REPLACEMENT THERAPY PARTICIPANTS AND NON-PARTICIPANTS .....	6-13
Fomenko Yu.V, Sukhostavets E, Hrechko N.B, Kuzina V.V, Mikhailenko N.M, Yaroslavska Yu.Yu, Skliar S.O, Mikulinska-Rudich Yu.M, Vlasov A.V, Smorodskiy V.O, Nazaryan R.S. PECULIARITIES OF THE SECOND MESIOBUCCAL CANAL IN MAXILLARY FIRST MOLAR: A RETROSPECTIVE ANALYSIS.....	14-20
Chikhashvili E, Kristesashvili J, Urjumelashvili M. EFFECTIVENESS OF COMBINED SURGICAL AND HORMONAL THERAPY IN TREATMENT OF ENDOMETRIOMAS.....	21-29
Lazzat I. Zhussupbekova, Dinara A. Nurkina, Saule M. Sarkulova, Galiya T. Smailova, Kassymzhomart N. Zholamanov. ACUTE FORMS OF CORONARY ARTERY DISEASE IN THE NOSOLOGICAL STRUCTURE OF HOSPITALIZATION OF YOUNG PEOPLE IN ALMATY CITY CARDIOLOGY CENTER.....	30-36
Alwashmi Emad, Alharbi Adel H, Almadi Abdulaziz S, Alhuraysi Abdulaziz, Almuhan Mousa M, Alharbi Badr. NOCTURNAL ENURESIS SYMPTOMS AND RISK FACTORS AMONG CHILDREN AND ADOLESCENTS IN QASSIM REGION, SAUDIARABIA.....	37-44
Askar Zh. Akhmetov, Tolkyn A. Bulegenov, Meirbek Zh. Aimagambetov, Nazarbek B. Omarov, Altay A. Dyusupov, Assel Zh. Baybussinova, Aldiyar E. Masalov, Samatbek T. Abdrakhmanov, Medet Ə. Ayenov. STATE OF INPATIENT MEDICAL CARE PATIENTS WITH ACUTE PANCREATITIS.....	45-51
Saad H . Abood, Liwaa A. Shihab, Ghufuran H. Abed, Thanon Y. Azzawi, Ahmed S. Abood. DETECTION OF MECA AND NUC GENES OF MULTI-DRUG RESISTANT STAPHYLOCOCCUS AUREUS ISOLATED FROM DIFFERENT CLINICAL SAMPLES.....	52-54
Sergey A. Apryatin, Vyacheslav I. Moiseenko, Raul R. Gainetdinov, Vera A. Apryatina. THE EFFECT OF INTRANASAL ADMINISTRATION OF BIOLOGICALLY ACTIVE SUBSTANCES OF AMINO ACID AND PEPTIDE NATURE ON THE MONOAMINE SYSTEMS OF THE BRAIN.....	55-67
Tchernev G, Broshtilova V, Kordeva S. DERMATOFIBROSARCOMA PROTUBERANS: WIDE LOCAL EXCISION AS DERMATOSURGICAL APPROACH WITH FAVOURABLE FINAL OUTCOME-CASE PRESENTATION AND SHORT UPDATE ON THERAPEUTIC OPTIONS.....	68-71
Yuuka Matsumoto, Takuma Hayashi, Yasuaki Amano, Kaoru Abiko, Ikuo Konishi. DEVELOPMENT OF ENDOSALPINGIOSIS IN PATIENTS WITH A HISTORY OF BREAST CANCER.....	72-76
Ilenko-Lobach N.V, Boychenko O.M, Ilenko N.M, Salomatina S.O, Nikolishyna E.V, Karnauh M.M, Voloshyna A.V, Zaitsev A.V. POSSIBILITY OF IMPROVING DISEASE PREDICTION USING MATHEMATICAL MODELS.....	77-79
Khabadze Z.S, Mer I.Ya, Fokina S.A, Mityushkina T.A, Kakabadze E.M, Badalov F.V, Dolzhikov N.A, Saeidyan S, Umarov A.Yu, Wehbe A. PROSPECTS AND LONG-TERM RESULTS AFTER ENDODONTIC SURGERY.....	80-86
Khatuna Kudava. NEVI IN CHILDREN: CLINICO-DERMOSCOPIC CONCEPTS ASSOCIATED WITH LOCATION.....	87-90
Jonathan Borges, Rashmi Aithmia, Jahnvi Mittal, Tarang Bhatnagar, Shivangi Gupta, Bhavuk Samrat. BREAST CANCER AND DIAGNOSTIC METHODS: UNDERSTANDING THE ROLE OF BRCA1 AND BRCA2.....	91-98
Kovaleva Kristina, Zulfiya Kachiyeva, Aigulim Abetova, Natalia Raspopova. GENETIC VARIANTS IN ANTIPSYCHOTIC METABOLISM: POLYMORPHISM PROFILES IN KAZAKH COHORT WITH PARANOID SCHIZOPHRENIA.....	99-103
Vakhtang Khelashvili, Tengiz Shiryaev, Omar Gogia. PERCUTANEOUS OCCLUSION OF MAJOR AORTOPULMONARY COLLATERALS IN TRANSPOSITION OF THE GREAT ARTERIES USING AMPLATZER PICCOLO OCCLUDERS: CASE REPORT.....	104-116
Ia Kusradze, Olia Rcheulishvili, Natia Karumidze, Sophio Rigvava, Aleksandre Rcheulishvili, Rusudan Goliadze, Luka Kamashidze, Alikyia Chipurupalli, Nunu Metreveli, Marine Goderdzishvili. PHAGE-BACTERIA INTERACTIONS UNDER METAL STRESS: A STUDY OF THE NOVEL STENOTROPHOMONAS MALTOPHILIA PHAGE VB_STM18.....	117-122
M.E. Azizova. PATHOMORPHOLOGICAL AND CLINICAL CHARACTERISTICS OF THE UTERUS IN COMBINED ADENOMYOSIS AND MYOMA.....	123-127
Grigoli Dzodzuashvili, Nino Chichua, Vladimer Margvelashvili, Giuli Margishvili, Natia Dzodzuashvili. STUDY OF ORAL HEALTH AND SUPPORTIVE STRUCTURES FOR PROSTHETIC RESTORATIONS IN METHADONE MAINTENANCE THERAPY BENEFICIARIES AND DRUG USERS.....	128-133

Noori Taha Alkhafaji, Mareb H. Ahmed, Bashar Rasim Karem. THE EFFECT OF VITAMIN D ON THE HISTOLOGICAL STRUCTURE OF LIVER AND LUNG IN MICE TREATED WITH AMPHOTERICIN B.....	134-141
Muratbekova Svetlana, Beth L. Leonberg, Kulbayeva Shynar, Duisenbina Zhanbota, Lissitsyn Yuriy. ASSESSING THE KNOWLEDGE LEVEL AND ATTITUDE TOWARDS PROVIDING NUTRITION CARE OF MEDICAL STUDENTS IN THE AKMOLA REGION OF THE REPUBLIC OF KAZAKHSTAN.....	142-147
Aldiyar E. Masalov, Meirbek Zh. Aimagambetov, Medet A. Auyenov, Samatbek T. Abdrakhmanov, Nazarbek B. Omarov, Altay A. Dyusupov, Tolkyn A. Bulegenov, Askar Zh. Akhmetov. IMPROVEMENT OF SURGICAL TREATMENT OF ACUTE BILIARY PANCREATITIS.....	148-155
Khabadze Z.S, Inozemtseva K.S, Bakaev Yu.A, Magomedov O.I, Kakabadze E.M, Badalov F.V, Saeidyan S, Umarov A.Yu, Wehbe A. A MODERN VIEW ON THE TREATMENT OF CLASS IV RECESSON ACCORDING TO MILLER.....	156-162
Christina Ejibishvili, Merab Kiladze, Ioseb Begashvili, George Grigolia. CORRELATION BETWEEN EJECTION FRACTION (EF) AND CORONARY SINUS BLOOD FLOW (CSBF) DURING OFF-PUMP CORONARY ARTERY BYPASS GRAFTING SURGERY.....	163-166
Tchernev G, Broshtilova V, Kordeva S. MULTIPLE MUSHROOM-LIKE GROWING CYLINDROMAS OF THE SCALP (TURBAN TUMOR) IN A PATIENT WITH BROOKE-SPIEGLER SYNDROME: UNIQUE MANIFESTATION IN A BULGARIAN PATIENT.....	167-170
Arnab Sain, Jack Song Chia, Nauman Manzoor, Minaal Ahmed Malik, Nadine Khayyat, Hamdoon Asim, Ahmed Elkilany, Otto Russell, Venera Derguti, Michele Halasa, Anushka Jindal, Fahad Hussain, Kanishka Wattage, Hoosai Manyar, Justin Wilson, Lulu Chamayi, Hannah Burton, Ansab Mahmood, Wilam Ivanga Alfred, Vivek Deshmukh, Abhinandan Kotian, Zain Sohail. BENNETT'S FRACTURE: A NARRATIVE REVIEW OF CURRENT LITERATURE.....	171-173
F. Kh. Umarov, J. J. Samatov. EARLY PREDICTORS OF NON-UNION OF DIAPHYSEAL TIBIAL FRACTURES BASED ON SCORING SYSTEMS.....	174-183
Satyanarayana Kummari, Aniket Madhukar Zope, Prachi Juyal, Pratibha Sharma, Sidhant Das, Sharin Koshy Varghese. DEEP LEARNING-BASED FRAMEWORK TO DETERMINE THE DEGREE OF COVID-19 INFECTIONS FROM CHEST X-RAY.....	184-187
Maghlakelidze Natalia, Zueva Marina V, Petriashvili Giorgi, Skliarenko Sofio. BINOCULAR INTERACTION IN AMBLYOPIA.....	188-191
Mariela Gaïbor-González, Diego Bonilla-Jurado, Ember Zumba-Novay, Cesar Guevara. STRATEGIC QUALITY MANAGEMENT OF PROCESSES IN NURSING SERVICES WITHIN INTERNAL AND GENERAL MEDICINE UNITS FOR A SUSTAINABLE FUTURE IN HEALTH SYSTEMS.....	192-200
Nugesha Grigalashvili, Lali Pkhaladze, Archil Khomasuridze. INTEGRATED MANAGEMENT OF OVARIAN ENDOMETRIOMAS: PRE- AND POST-SURGICAL USE OF DIENOGEST.....	201-205
S. Rigvava, I Kusradze, N. Karumidze, M. Chichashvili, I. Tchgkonია, M. Goderdzishvili. SMALL BUT MIGHTY: CHARACTERIZATION OF VB_SPY_7, A LYTIC PHAGE TARGETING STREPTOCOCCUS PYOGENES.....	206-210
Gorbik E.V, Ohurtsov O.S, Heranin S.I, Kolba O.O, Breslavets N.M, Sazonova O.M, Kysylenko K.V, Alekseeva V.V. ANATOMY OF THE MAXILLARY SINUS: IMPLICATIONS FOR ODONTOGENIC SINUSITIS DEVELOPMENT.....	211-216
Zviad Kereselidze, Lela Kopaleishvili, Kakha Nadaraia, Kakhaber Chelidze, Vakhtang Chumburize. CARVEDILOL IN PATIENTS WITH UNCONTROLLED AND RESISTANT ARTERIAL HYPERTENSION.....	217-224
Mirian Getsadze, Sofia Chedia. STUDY OF ORBITAL NEOPLASMS BY MAGNETIC RESONANCE IMAGING PROCEDURE.....	225-233

## POSSIBILITY OF IMPROVING DISEASE PREDICTION USING MATHEMATICAL MODELS

Ilenko-Lobach N.V<sup>1</sup>, Boychenko O.M<sup>1</sup>, Ilenko N.M<sup>1</sup>, Salomatina S.O<sup>2</sup>, Nikolishyna E.V<sup>1</sup>, Karnauh M.M<sup>3</sup>, Voloshyna A.V<sup>3</sup>, Zaitsev A.V<sup>1</sup>.

<sup>1</sup>Associate Professor of the Department of Therapeutic Stomatology, Poltava State Medical University, Poltava, Ukraine.

<sup>2</sup>Associate Professor of the Department of Therapeutic Stomatology, Dnipro State Medical University, Ukraine.

<sup>3</sup>Associate Professor of the Department of Therapeutic Stomatology, Zaporizhia State Medical and Pharmaceutical University, Ukraine.

### Abstract.

Throughout their existence, people have sought to know what tomorrow holds for them. It was especially important to learn about a full life, illness or death. Science is the essence of human knowledge, which is organized according to certain principles, a real connection of judgments, predictions and problems of reality and its individual spheres or aspects.

Various indexes are used in modern periodontology. Index assessment allows you to succinctly and conveniently display the static state of periodontal tissues in quantitative terms. With their help, you can get an idea of the prevalence, degree of severity, the course of the inflammatory-destructive or dystrophic process, evaluate the effectiveness of the treatment and the quality of dispensary work in a specific patient or dispensary group. Among the indices, clinical ones are especially defined. They are the main group of indices that allows the clinician to assess the state of the periodontal tissues in a variety of ways. With their help, you can get a general idea of the nature of the course of the disease, plan the amount of medical interventions, evaluate the effectiveness of the treatment and dispensation.

Various indexes are used in modern periodontology. Index assessment allows you to succinctly and conveniently display the static state of periodontal tissues in quantitative terms. With their help, you can get an idea of the prevalence, degree of severity, the course of the inflammatory-destructive or dystrophic process, evaluate the effectiveness of the treatment and the quality of dispensary work in a specific patient or dispensary group. Among the indices, clinical ones are especially defined. They are the main group of indices that allows the clinician to assess the state of the periodontal tissues in a variety of ways. With their help, you can get a general idea of the nature of the course of the disease, plan the amount of medical interventions, evaluate the effectiveness of the treatment and dispensation.

**Key words.** Forecasting, mathematical modeling, periodontal disease.

### Introduction.

Throughout their existence, people have sought to know what tomorrow holds for them. It was especially important to learn about a full life, illness or death. Science is the essence of human knowledge, which is organized according to certain principles, a real connection of judgments, predictions and problems of reality and its individual spheres or aspects. The growth of human experience is ensured through scientific works. Scientific forecasting is a type of theoretical activity that consists in defining and describing certain phenomena of nature, social life, mental states that are absent or not known now, but may arise or be studied and discovered in the future. The problem of scientific forecasting is relevant for many

spheres of life, including medicine. [1]. Currently, in the time of evidence-based medicine and the information and legal space, patients have high demands on the results of treatment, which forces doctors to clearly predict these results. Unfortunately, the existing forecasting methods and tools do not always sufficiently predict the course of the disease, remain complex and not integrated into practical activities [2.].

Mathematical modeling as a method of scientific knowledge began to be used by mankind many centuries ago, from the moment when the foundations of differential and integral calculus were laid. But its use in biomedical fields began only in the 80s of the 19th century. Modeling in medicine allows establishing deeper and more complex relationships between theory and experiment. A whole series of studies is impossible without it. In addition, some experiments are harmful to health. With the help of modeling, a number of different models can be developed on one set of information. You can also make your additions to the model. Thanks to modeling in the second half of the 20th century. immunology began to develop rapidly [3].

Many calculations are performed on computers using programs designed for this purpose. One of them is Microsoft Office Excel. With its help, you can simplify many tasks of any specialist who uses a computer in his work. This mathematical processor has been used in the field of data analysis for decades and has become an indispensable component of every computer. Excel provides users with the following capabilities:

1. Allows you to create dynamic and easy-to-manage tables.
2. Allows you to perform various calculations and operations on data. Formulas can include arithmetic operations, functions, conditional expressions, and more. It allows you to automate calculations and perform complex analytical tasks.
3. Provides ample opportunities to create various types of graphs and charts, facilitating effective analysis.
4. Allows you to filter and sort large amounts of information.
5. Excel has built-in analytical features that help you perform various types of analysis, including regression analysis, correlations, trends, and more.
6. Allows you to use predictive models to determine future trends.

To work with this program, the Statsoft company also developed the "STATISTICA" system, which is one of the most popular statistical programs for finding patterns, forecasting, classification, and visualization of material. It is used in various spheres of human activity, in particular in medicine [4].

The listed opportunities create incredible conditions for conducting various researches, which is very important for the advancement of human knowledge. Therefore, it is of interest to know how well modern scientists in medicine use the listed computer support for prediction. This is the relevance of our work.



**The purpose of the work:** Analysis of modern clinical research such as forecasting using the capabilities of the computer program Excel, in particular in dental theory and practice.

**The task:** To evaluate the peculiarities of the use of the Excel device in predicting treatment in modern dental research and to make further assumptions regarding the improvement of this practice.

**Scientific novelty:** The results of the assessment of the features of the use of the Excel device for predicting treatment in modern dental research were obtained, and further assumptions were made regarding the improvement of this practice.

**Practical significance:** The conducted analysis has theoretical and practical significance for therapeutic stomatology.

## Materials and Methods.

The materials were the results of clinical studies conducted at the dental center of the National Medical University named after O.O. Bogomolets; the results of a scientific study conducted in the cardiology department of the Lviv Emergency Medical Care Hospital and the Dental Medical Center of Danylo Halytsky Lviv National Medical University; the results of a scientific study conducted in the rheumatology department of the ENT hospital "Lviv Regional Clinical Hospital" and at the Department of Therapeutic Stomatology, Periodontology and Stomatology of the Danylo Halytsky Lviv National Medical University.

Analysis of the aforementioned clinical studies. Mathematical analysis based on elements of regression analysis - approximation, which was carried out using the computer program Excel, which we used repeatedly [6], as well as other capabilities of this program [7].

## Results.

Various indexes are used in modern periodontology. Index assessment allows you to succinctly and conveniently display the static state of periodontal tissues in quantitative terms. With their help, you can get an idea of the prevalence, degree of severity, the course of the inflammatory-destructive or dystrophic process, evaluate the effectiveness of the treatment and the quality of dispensary work in a specific patient or dispensary group. Among the indices, clinical ones are especially defined. They are the main group of indices that allows the clinician to assess the state of the periodontal tissues in a variety of ways. With their help, you can get a general idea of the nature of the course of the disease, plan the amount of medical interventions, evaluate the effectiveness of the treatment and dispensation.

The PMA (papillary-marginal-alveolar (cellular)) index belongs to the indices that determine the degree of inflammation of the gums. The index was proposed by I. Schour and M. Massler in 1948. It is one of the most common - the first basic indices of gingival inflammation, which was subsequently modified by various authors. The modification of C. Parma in 1960 was the most convenient [8-10]. The index allows you to objectively assess the degree of inflammation of the gums and is still used in the C. Parma modification [11,12]. Such long-term viability of this clinical method draws attention to it, makes it more closely familiar with the example of its use in the sense of forecasting the treatment of periodontal diseases, and possibly another embodiment.

It is interesting to see how well the treatment indicators will be displayed with the help of PMA when the mathematical apparatus of the Excel computer program is applied to it. This method was used by us regarding the relationship between representatives of the oral microbiome and proved good results [8]. The mathematical apparatus is a universal method of studying a real phenomenon or process. Therefore, we tried to involve this concept in identifying the possibilities of predicting the condition of periodontal tissues using an index assessment.

To find out the effectiveness of the application of different types of scaling at the first stage of treatment of periodontal diseases in 2017 at the dental center of the National Medical University of O.O. Bogomolets a study was conducted.

Patients with equivalent periodontal conditions (chronic generalized periodontitis, chronic catarrhal gingivitis of mild and moderate severity, chronic hypertrophic gingivitis of mild severity) were divided into groups according to the types of scaling:

Patients with equivalent periodontal conditions (chronic generalized periodontitis, chronic catarrhal gingivitis of mild and moderate severity, chronic hypertrophic gingivitis of mild severity) were divided into groups according to the type of scaling: using instrumental scaling (using hand tools), using electromechanical scaling (using an ultrasonic tool) and using combined scaling (using ultrasound and hand tools). They underwent a clinical examination, which included an index assessment of the level of periodontal inflammation using the PMA index (Table 1) [9].


**Table 1.** Changes in indicators of the PMA index according to C. Parma (1960), % in patients using different types of scaling ( $M \pm m$ ).

Types of scaling	Research deadlines			
	1 day	2 day	14 day	after 3 months
Instrumental	24,30±1,14	26,01±1,15	18,33±1,15	20,30±1,14
Electromechanical	23,50±1,11	15,13±1,15	10,65±1,15	21,29±1,11
Combined	23,18±1,11	20,15±1,15	4,91±1,15	14,58±1,11


On the basis of the obtained results, conclusions were drawn regarding the advantage of the combined method of removing dental deposits - it combines the advantages of instrumental and ultrasonic types of scaling and allows you to avoid complications arising from the use of these methods separately.

We have created mathematical models of the dynamics of periodontal inflammation in the above-mentioned groups of patients - with the help of approximation, graphs of changes in the PMA index over 3 months have been created, graph trends and their mathematical formulas have been reproduced (Figure 1) [13].

Thus, for the group of patients who underwent instrumental scaling, a mathematical model with the best approximation reliability ( $R^2$ ) was created when determining the polynomial function according to the formula  $y = 0.0063 x^2 - 0.6306 x + 26.032$ .  $R^2$  was 0.9277. For the group of patients who underwent electromechanical scaling, a mathematical model was created with the best approximation reliability ( $R^2$ ) when determining the polynomial function according to the formula  $y = 0.0099 x^2 - 0.8887 x + 20.795$ .  $R^2$  was 0.7239. For the group of patients



**Figure 1.** Graphs, trends, and formulas for changes in PMA index indicators according to C. Parma (1960) (%) in patients using different types of scaling within 3 months.



**Figure 2.** Graphs of changes in indicators of the PMA index according to C. Parma (1960) (%) in patients using different types of scaling during 12 months.

who underwent combined scaling, a mathematical model was created with the best approximation reliability ( $R^2$ ) when determining the polynomial function according to the formula  $y = 0.0167x^2 - 1.6068x + 24.067$ .  $R^2$  was 0.9944.

After that, with the help of Excel, we created graphs that reproduced the dynamics of periodontal inflammation (according to the PMA index) over 12 months (Figure 2).

The resulting graphic models visualize how inflammation can develop in the periodontal tissues if nothing is involved in reducing its progression.

In 2022, a dissertation was defended, devoted to the solution of measures to optimize the treatment and prevention of chronic generalized periodontitis in patients with ischemic heart disease through the use of statin-based drugs. Patients with ischemic heart disease who were undergoing inpatient treatment at the cardiology department of the Emergency Medical Care Hospital of the city of Lviv and patients with a diagnosis of chronic generalized periodontitis without somatic diseases who applied for treatment at the Danylo Halytskyi Dental Medical Center of LNMU.

Three groups were formed from them: the main group and comparison group 1 - patients with ischemic heart disease and

chronic generalized periodontitis of the initial I and II degree of severity and comparison group 2 - patients with chronic generalized periodontitis, in which the presence of somatic pathology was not indicated in the anamnesis [14].

The results of determining the PMA index before treatment, after it and in the long term are shown in Table 2.

**Table 2.** Results of treatment and prevention of chronic generalized periodontitis according to the PMA index in the studied groups ( $M \pm m, \%$ ).

Research groups	Terms of research			
	Before treatment	After treatment 7 days	After treatment 1 month	After treatment 6 months
Main group	49,89	22,78	20,88	19,09
Comparison group 1	48,42	34,29	31,86	38,48
Comparison group 2	39,29	21,56	20,94	26,47

Based on the obtained results, the therapeutic effectiveness of the developed gel composition based on atorvastatin was proven

and it was recommended as an additional local medication in the complex treatment and prevention of inflammatory and dystrophic-inflammatory diseases of periodontal tissues in patients with ischemic heart disease.

We have created mathematical models of the dynamics of periodontal inflammation in the indicated groups of patients - by means of approximation, graphs of changes in the PMA index over 6 months were invented, graph trends and their mathematical formulas were reproduced (Figure 3).

The following indicators were invented:

1. For the main group, the best approximation reliability of  $R^2=0.6045$  was when determining the polynomial function with the following formula:  $y=0.0044x^2-0.9108x+41.313$ .

2. For comparison group 1, the best approximation reliability of  $R^2=0.6023$  was when determining the polynomial function with the following formula:  $y=0.0029x^2-0.5465x+44.196$ .

3. For comparison group 2, the best approximation reliability of  $R^2=0.497$  was when determining the polynomial function with the following formula:  $y=0.003x^2-0.5768x+33.61$ .

After that, with the help of Excel, we created graphs that reproduced the dynamics of periodontal inflammation (according to the PMA index) over 12 months (Figure 4).

The resulting graphic models visualize how inflammation can develop in the periodontal tissues if nothing is involved in reducing its progression.

In Lviv, in 2023, a dissertation work was defended, which presented a theoretical generalization and a new solution to an urgent task of modern dentistry - clinical and laboratory substantiation of the effectiveness of the prevention and treatment of periodontal diseases in patients with gout based on the results of clinical, biochemical, immunological and physicochemical blood tests and oral fluid.

In order to achieve the goal, two groups of patients were created: the main group with chronic generalized periodontitis (initial-I and II degrees of severity) was treated according to the

therapeutic and preventive scheme developed by the doctoral student: additionally prescribed the drugs "Monmorol" (ToV Nutrimed, Ukraine); "Biotrit-denta" (NPA Odesa Biotechnology, Ukraine); the drug "Acerola S-500" with bioflavonoids (ToV Nutrimed, Ukraine). In the control group, traditional methods were used for the treatment of periodontal diseases (gingivitis, chronic generalized periodontitis initial-I and II degrees of severity, periodontitis) in accordance with generally accepted protocols. The papillary-marginal-alveolar index (RMA, Parma, 1960) was used to assess the degree of inflammation of the gums. The proposed treatment and prevention complex made it possible to achieve remission of generalized periodontitis.


The results of determining the PMA index before treatment, after it and in the long term are shown in Table 3.

**Table 3.** Results of treatment and prevention of chronic generalized periodontitis according to the PMA index in the studied groups ( $M\pm m, \%$ ).


Terms of observation	Main group PMA, %	Control group PMA, %
Before treatment	58,30±10,31	56,89±10,14
1 month after treatment	33,21±3,16	42,88±6,32
6 months after treatment	31,09±3,14	44,19±6,53
12 months after treatment	40,83±5,55	58,02±10,74

We have created mathematical models of the dynamics of periodontal inflammation in the indicated groups of patients - by means of approximation, graphs of changes in the PMA index over 12 months were invented, graph trends and their mathematical formulas were reproduced (Figure 5).


For the main group, the best approximation reliability ( $R^2=0.6216$ ) was when defining the logarithmic function




**Figure 3.** Graphs, trends and formulas of changes in PMA index indicators according to C. Parma (1960) (%) in patients with chronic generalized periodontitis on the background of ischemic heart disease and without somatic pathology during 6 months.



**Figure 4.** Graphs of changes in PMA index according to C. Parma (1960) (%) in patients with chronic generalized periodontitis on the background of ischemic heart disease and without somatic pathology during 12 months.



**Figure 5.** Graphs, trends, and formulas for changes in PMA index indicators according to C. Parma (1960) (%) in patients with chronic generalized periodontitis on the background of gout within 12 months.



**Figure 6.** Graphs of changes in indicators of the PMA index according to C. Parma (1960) (%) in patients with gout during 36 months.

according to the following formula:  $y = -3.704 \ln(x) + 54.265$ . For the control group, the best approximation reliability ( $R^2 = 0.6864$ ) was when determining the polynomial function according to the following formula:  $y = 0.0004x^2 - 0.1276x + 52.375$ . We have created mathematical models of the dynamics of periodontal inflammation in the indicated groups of patients - by means of approximation, graphs of changes in the PMA index over 36 months were invented, graph trends and their mathematical formulas were reproduced (Figure 6).

### Discussion.

With the help of the Excel computer program, we obtained prognostic mathematical models of the disease when applying various treatment measures to its stabilization.

In the study of Dikova I.H. for the group of patients who underwent instrumental scaling, a mathematical model was created with the best approximation reliability ( $R^2 = 0.9277$ ) when determining the polynomial function according to the formula  $y = 0.0063 x^2 - 0.6306 x + 26.032$ . For the group

of patients who underwent electromechanical scaling, a mathematical model was created with the best approximation reliability ( $R^2=0.7239$ ) when determining the polynomial function according to the formula  $y = 0.0099 x^2 - 0.8887 x + 20.795$ .

For the group of patients who underwent combined scaling, a mathematical model with the best approximation reliability ( $R^2=0.9944$ ) was created when determining the polynomial function according to the formula  $y = 0.0167 x^2 - 1.6068 x + 24.067$ .

In the study of O.V. Skibchyk. The following indicators were invented:

1. For the main group, the best approximation reliability ( $R^2=0.6045$ ) was when determining the polynomial function according to the formula  $y=0.0044x^2-0.9108x+41.313$ .

2. For comparison group 1, the best approximation reliability ( $R^2=0.6023$ ) was when determining the polynomial function according to the formula  $y=0.0029x^2-0.5465x+44.196$ .

3. For comparison group 2, the best approximation reliability ( $R^2=0.497$ ) was when determining the polynomial function according to the formula  $y=0.003x^2-0.5768x+33.61$ .

In the study Gnid M.R. for the main group, the best approximation reliability ( $R^2=0.6216$ ) was when defining the logarithmic function according to the following formula:  $y=-3.704\ln(x)+54.265$ . For the control group, the best approximation reliability ( $R^2=0.6864$ ) is based on the following formula:  $y=0.0004x^2-0.1276x+52.375$ .

The reliability index can vary from 0 to 1. The higher it is, the more reliable the approximation.

It is believed that when the value of this indicator is 0.85 and above, the correspondence of the simulated function to reality can be considered the most reliable.

In essence, all invented mathematical models, except for one, are formulas of a function of the second degree (quadratic). But in the case of research Hnid M.R. for the main group, the best approximation reliability was found when defining the logarithmic function.

In the study of Dikova I.G. according to the mathematical models created by us, a stable remission is achieved within three months. After 100 days, the progression of the disease begins. In addition, the combined method of scaling after the end of remission gives the fastest further progression of the disease.

In the study of O.V. Skibchyk. according to the mathematical models created by us, a stable remission is also achieved within three months. Then there is a slow progression of the disease, in which the main group is the first to reach one hundred percent.

In the study Hnid M.R. according to the mathematical models created by us, a stable remission is observed for 200 days. After that, the progression of the disease begins in the comparison group. And in the main group, on the contrary, remission goes to the stabilization phase.

From July 1, 2018, the Procedure for the provision of primary medical care entered into force, and the dispensary became a thing of the past. In its place, scheduled screenings have been introduced, the conduct of which is determined by the order of the Ministry of Health of Ukraine dated March 19, 2018 No. 504 "On approval of the Procedure for providing primary medical care".

## Conclusion.

Modelling of the disease process is possible using the tools of the mathematical processor Microsoft Office Excel. But in our case, the obtained mathematical models do not always have the appropriate degree of reliability. This indicates the need to refine the mathematical apparatus due to the databases used in medical research. Reproduction of mathematical models with different formulas can indicate not only the imperfection of the used approach. It can also be a manifestation of the intervention of therapeutic agents, which were used to eliminate or stabilize the disease, in the work of other body systems besides the dental and jaw system. Or something else, which can also be caused by the vital activity of the human body or the influence of the surrounding environment on it. In our study, the case of inconsistency of one mathematical model with others has positive consequences.

The different period of remission also attracts attention. In connection with the introduction of the screening assessment, the modelling of the further development of the disease is gaining relevance. This indicates that doctors in their research should use the construction of mathematical models both in the sense of predicting the consequences of treatment and the progress of the disease, as well as to determine the date of a repeat visit to the doctor and to study the essence of the pathology itself. This can be beneficial in understanding the studied disease state, as well as in creating new approaches to its elimination and disease prevention (correction).

**Prospects for further research:** are to clarify the possibility of improving the quality of forecasting the consequences of treatment and the progress of the disease using mathematical modeling in dental practice.

**Conflict of interest.** The authors declare no conflict of interest.

## REFERENCES

1. Birta G.O, Burgu Y.G. Methodology and organization of scientific research: teaching manual. "Center for Educational Literature". 2014:142.
2. V'yun G.I. Forecasting the results of periodontal treatment of patients with generalized periodontitis. *Modern Dentistry*. 2018;3:32-35.
3. Melnyk M. A, Danylychuk O. M. Mathematical model. Application of higher mathematics in the process of training specialists in the fields of medicine and biology. *Bulletin of the Student Scientific Society of the Vasyl Stus Donetsk National University*. 2021;1:280-284.
4. Klymchuk V.O. Mathematical methods in psychology. Textbook for students of psychological specialties. — K.: *Osvita Ukrainy*. 2009:288.
5. Zaitsev A. V, Boychenko O. M, Kotelevska N. V, et al. Method of determining indicators connection of the state of periodontal inflammation and oral microbiota. *Bulletin of problems in biology and medicine*. 2023;168:270-275.
6. Excel 2013–2016: textbook / Compiled by: Dyachuk S.F. — Ternopil: Ivan Pulyua TNTU Publishing House. 2021:308.
7. Boiko DI, Skrypnykov AM, Shkodina AD, et al. Circadian rhythm disorder and anxiety as mental health complications in post-COVID-19. *Environ Sci Pollut Res Int*. 2022;29:28062-28069.

8. Dikova I.G, Gorodenko E.O. Justification of the prevention of complications when using different types of scaling. *Modern Dentistry*. 2017:22-24.
10. Kutsevlyak V.F. Index assessment of periodontal status. V.F. Kutsevlyak, Yu.V. Lakhtin: textbook. 2nd ed. revised and supplemented. - Sumy: Publishing and Production Enterprise "Mriya". 2015:104.
11. Bezvushko E.V, Malko N.V. Correlation between papillary-marginal-alveolar index and immunological parameters of oral fluid in children with chronic catarrhal gingivitis. *Bulletin of Problems in Biology and Medicine: Ukrainian Scientific and Practical Journal*. 2023;4:413-417.
12. Errors in constructing graphs in Excel – YouTube.2024. <https://www.youtube.com/watch?v=DddvuwR8rqU>.
13. Skybchik OV. Optimization of prevention and complex treatment of generalized periodontitis with the use of statins in patients with ischemic heart disease: dissertation of doctor of philosophy: Skybchik Oksana Vasylivna. Lviv. 2022:258.
14. Errors in plotting graphs in Excel – YouTube. <https://www.youtube.com/watch?v=DddvuwR8rqU>. 2024