

GEORGIAN MEDICAL NEWS

ISSN 1512-0112

NO 2 (359) Февраль 2025

ТБИЛИСИ - NEW YORK



ЕЖЕМЕСЯЧНЫЙ НАУЧНЫЙ ЖУРНАЛ

Медицинские новости Грузии
საქართველოს სამედიცინო სიახლენი

GEORGIAN MEDICAL NEWS

Monthly Georgia-US joint scientific journal published both in electronic and paper formats of the Agency of Medical Information of the Georgian Association of Business Press.
Published since 1994. Distributed in NIS, EU and USA.

GMN: Georgian Medical News is peer-reviewed, published monthly journal committed to promoting the science and art of medicine and the betterment of public health, published by the GMN Editorial Board since 1994. GMN carries original scientific articles on medicine, biology and pharmacy, which are of experimental, theoretical and practical character; publishes original research, reviews, commentaries, editorials, essays, medical news, and correspondence in English and Russian.

GMN is indexed in MEDLINE, SCOPUS, PubMed and VINITI Russian Academy of Sciences. The full text content is available through EBSCO databases.

GMN: Медицинские новости Грузии - ежемесячный рецензируемый научный журнал, издаётся Редакционной коллегией с 1994 года на русском и английском языках в целях поддержки медицинской науки и улучшения здравоохранения. В журнале публикуются оригинальные научные статьи в области медицины, биологии и фармации, статьи обзорного характера, научные сообщения, новости медицины и здравоохранения. Журнал индексируется в MEDLINE, отражён в базе данных SCOPUS, PubMed и ВИНТИ РАН. Полнотекстовые статьи журнала доступны через БД EBSCO.

GMN: Georgian Medical News – საქართველოს სამედიცინო სიახლენი – არის ყოველთვიური სამეცნიერო სამედიცინო რეცენზირებადი ჟურნალი, გამოიცემა 1994 წლიდან, წარმოადგენს სარედაქციო კოლეგიისა და აშშ-ის მეცნიერების, განათლების, ინდუსტრიის, ხელოვნებისა და ბუნებისმეტყველების საერთაშორისო აკადემიის ერთობლივ გამოცემას. GMN-ში რუსულ და ინგლისურ ენებზე ქვეყნდება ექსპერიმენტული, თეორიული და პრაქტიკული ხასიათის ორიგინალური სამეცნიერო სტატიები მედიცინის, ბიოლოგიისა და ფარმაციის სფეროში, მიმოხილვითი ხასიათის სტატიები.

ჟურნალი ინდექსირებულია MEDLINE-ის საერთაშორისო სისტემაში, ასახულია SCOPUS-ის, PubMed-ის და ВИНТИ РАН-ის მონაცემთა ბაზებში. სტატიების სრული ტექსტი ხელმისაწვდომია EBSCO-ს მონაცემთა ბაზებიდან.

WEBSITE

www.geomednews.com

К СВЕДЕНИЮ АВТОРОВ!

При направлении статьи в редакцию необходимо соблюдать следующие правила:

1. Статья должна быть представлена в двух экземплярах, на русском или английском языках, напечатанная через **полтора интервала на одной стороне стандартного листа с шириной левого поля в три сантиметра**. Используемый компьютерный шрифт для текста на русском и английском языках - **Times New Roman (Кириллица)**, для текста на грузинском языке следует использовать **AcadNusx**. Размер шрифта - **12**. К рукописи, напечатанной на компьютере, должен быть приложен CD со статьей.

2. Размер статьи должен быть не менее десяти и не более двадцати страниц машинописи, включая указатель литературы и резюме на английском, русском и грузинском языках.

3. В статье должны быть освещены актуальность данного материала, методы и результаты исследования и их обсуждение.

При представлении в печать научных экспериментальных работ авторы должны указывать вид и количество экспериментальных животных, применявшиеся методы обезболивания и усыпления (в ходе острых опытов).

4. К статье должны быть приложены краткое (на полстраницы) резюме на английском, русском и грузинском языках (включающее следующие разделы: цель исследования, материал и методы, результаты и заключение) и список ключевых слов (key words).

5. Таблицы необходимо представлять в печатной форме. Фотокопии не принимаются. **Все цифровые, итоговые и процентные данные в таблицах должны соответствовать таковым в тексте статьи**. Таблицы и графики должны быть озаглавлены.

6. Фотографии должны быть контрастными, фотокопии с рентгенограмм - в позитивном изображении. Рисунки, чертежи и диаграммы следует озаглавить, пронумеровать и вставить в соответствующее место текста **в tiff формате**.

В подписях к микрофотографиям следует указывать степень увеличения через окуляр или объектив и метод окраски или импрегнации срезов.

7. Фамилии отечественных авторов приводятся в оригинальной транскрипции.

8. При оформлении и направлении статей в журнал МНГ просим авторов соблюдать правила, изложенные в «Единых требованиях к рукописям, представляемым в биомедицинские журналы», принятых Международным комитетом редакторов медицинских журналов - <http://www.spinesurgery.ru/files/publish.pdf> и http://www.nlm.nih.gov/bsd/uniform_requirements.html В конце каждой оригинальной статьи приводится библиографический список. В список литературы включаются все материалы, на которые имеются ссылки в тексте. Список составляется в алфавитном порядке и нумеруется. Литературный источник приводится на языке оригинала. В списке литературы сначала приводятся работы, написанные знаками грузинского алфавита, затем кириллицей и латиницей. Ссылки на цитируемые работы в тексте статьи даются в квадратных скобках в виде номера, соответствующего номеру данной работы в списке литературы. Большинство цитированных источников должны быть за последние 5-7 лет.

9. Для получения права на публикацию статья должна иметь от руководителя работы или учреждения визу и сопроводительное отношение, написанные или напечатанные на бланке и заверенные подписью и печатью.

10. В конце статьи должны быть подписи всех авторов, полностью приведены их фамилии, имена и отчества, указаны служебный и домашний номера телефонов и адреса или иные координаты. Количество авторов (соавторов) не должно превышать пяти человек.

11. Редакция оставляет за собой право сокращать и исправлять статьи. Корректур авторам не высылаются, вся работа и сверка проводится по авторскому оригиналу.

12. Недопустимо направление в редакцию работ, представленных к печати в иных издательствах или опубликованных в других изданиях.

При нарушении указанных правил статьи не рассматриваются.

REQUIREMENTS

Please note, materials submitted to the Editorial Office Staff are supposed to meet the following requirements:

1. Articles must be provided with a double copy, in English or Russian languages and typed or computer-printed on a single side of standard typing paper, with the left margin of 3 centimeters width, and 1.5 spacing between the lines, typeface - **Times New Roman (Cyrillic)**, print size - 12 (referring to Georgian and Russian materials). With computer-printed texts please enclose a CD carrying the same file titled with Latin symbols.

2. Size of the article, including index and resume in English, Russian and Georgian languages must be at least 10 pages and not exceed the limit of 20 pages of typed or computer-printed text.

3. Submitted material must include a coverage of a topical subject, research methods, results, and review.

Authors of the scientific-research works must indicate the number of experimental biological species drawn in, list the employed methods of anesthetization and soporific means used during acute tests.

4. Articles must have a short (half page) abstract in English, Russian and Georgian (including the following sections: aim of study, material and methods, results and conclusions) and a list of key words.

5. Tables must be presented in an original typed or computer-printed form, instead of a photocopied version. **Numbers, totals, percentile data on the tables must coincide with those in the texts of the articles.** Tables and graphs must be headed.

6. Photographs are required to be contrasted and must be submitted with doubles. Please number each photograph with a pencil on its back, indicate author's name, title of the article (short version), and mark out its top and bottom parts. Drawings must be accurate, drafts and diagrams drawn in Indian ink (or black ink). Photocopies of the X-ray photographs must be presented in a positive image in **tiff format**.

Accurately numbered subtitles for each illustration must be listed on a separate sheet of paper. In the subtitles for the microphotographs please indicate the ocular and objective lens magnification power, method of coloring or impregnation of the microscopic sections (preparations).

7. Please indicate last names, first and middle initials of the native authors, present names and initials of the foreign authors in the transcription of the original language, enclose in parenthesis corresponding number under which the author is listed in the reference materials.

8. Please follow guidance offered to authors by The International Committee of Medical Journal Editors guidance in its Uniform Requirements for Manuscripts Submitted to Biomedical Journals publication available online at: http://www.nlm.nih.gov/bsd/uniform_requirements.html
http://www.icmje.org/urm_full.pdf

In GMN style for each work cited in the text, a bibliographic reference is given, and this is located at the end of the article under the title "References". All references cited in the text must be listed. The list of references should be arranged alphabetically and then numbered. References are numbered in the text [numbers in square brackets] and in the reference list and numbers are repeated throughout the text as needed. The bibliographic description is given in the language of publication (citations in Georgian script are followed by Cyrillic and Latin).

9. To obtain the rights of publication articles must be accompanied by a visa from the project instructor or the establishment, where the work has been performed, and a reference letter, both written or typed on a special signed form, certified by a stamp or a seal.

10. Articles must be signed by all of the authors at the end, and they must be provided with a list of full names, office and home phone numbers and addresses or other non-office locations where the authors could be reached. The number of the authors (co-authors) must not exceed the limit of 5 people.

11. Editorial Staff reserves the rights to cut down in size and correct the articles. Proof-sheets are not sent out to the authors. The entire editorial and collation work is performed according to the author's original text.

12. Sending in the works that have already been assigned to the press by other Editorial Staffs or have been printed by other publishers is not permissible.

**Articles that Fail to Meet the Aforementioned
Requirements are not Assigned to be Reviewed.**

ავტორთა საქურაღებოლ!

რედაქციაში სტატიის წარმოდგენისას საჭიროა დაიცვათ შემდეგი წესები:

1. სტატია უნდა წარმოადგინოთ 2 ცალად, რუსულ ან ინგლისურ ენებზე დაბეჭდილი სტანდარტული ფურცლის 1 გვერდზე, 3 სმ სიგანის მარცხენა ველისა და სტრიქონებს შორის 1,5 ინტერვალის დაცვით. გამოყენებული კომპიუტერული შრიფტი რუსულ და ინგლისურენოვან ტექსტებში - **Times New Roman (Кириллица)**, ხოლო ქართულენოვან ტექსტში საჭიროა გამოვიყენოთ **AcadNusx**. შრიფტის ზომა – 12. სტატიას თან უნდა ახლდეს CD სტატიით.

2. სტატიის მოცულობა არ უნდა შეადგენდეს 10 გვერდზე ნაკლებს და 20 გვერდზე მეტს ლიტერატურის სიის და რეზიუმეების (ინგლისურ, რუსულ და ქართულ ენებზე) ჩათვლით.

3. სტატიაში საჭიროა გაშუქდეს: საკითხის აქტუალობა; კვლევის მიზანი; საკვლევი მასალა და გამოყენებული მეთოდები; მიღებული შედეგები და მათი განსჯა. ექსპერიმენტული ხასიათის სტატიების წარმოდგენისას ავტორებმა უნდა მიუთითონ საექსპერიმენტო ცხოველების სახეობა და რაოდენობა; გაუტკივარებისა და დაძინების მეთოდები (მწვავე ცდების პირობებში).

4. სტატიას თან უნდა ახლდეს რეზიუმე ინგლისურ, რუსულ და ქართულ ენებზე არანაკლებ ნახევარი გვერდის მოცულობისა (სათაურის, ავტორების, დაწესებულების მითითებით და უნდა შეიცავდეს შემდეგ განყოფილებებს: მიზანი, მასალა და მეთოდები, შედეგები და დასკვნები; ტექსტუალური ნაწილი არ უნდა იყოს 15 სტრიქონზე ნაკლები) და საკვანძო სიტყვების ჩამონათვალი (key words).

5. ცხრილები საჭიროა წარმოადგინოთ ნაბეჭდი სახით. ყველა ციფრული, შემაჯამებელი და პროცენტული მონაცემები უნდა შეესაბამებოდეს ტექსტში მოყვანილს.

6. ფოტოსურათები უნდა იყოს კონტრასტული; სურათები, ნახაზები, დიაგრამები - დასათაურებული, დანომრილი და სათანადო ადგილას ჩასმული. რენტგენოგრამების ფოტოასლები წარმოადგინეთ პოზიტიური გამოსახულებით **tiff** ფორმატში. მიკროფოტოსურათების წარწერებში საჭიროა მიუთითოთ ოკულარის ან ობიექტივის საშუალებით გადიდების ხარისხი, ანათალების შედეგის ან იმპრეგნაციის მეთოდი და აღნიშნოთ სურათის ზედა და ქვედა ნაწილები.

7. სამამულო ავტორების გვარები სტატიაში აღინიშნება ინიციალების თანდართვით, უცხოურისა – უცხოური ტრანსკრიპციით.

8. სტატიას თან უნდა ახლდეს ავტორის მიერ გამოყენებული სამამულო და უცხოური შრომების ბიბლიოგრაფიული სია (ბოლო 5-8 წლის სიღრმით). ანბანური წყობით წარმოდგენილ ბიბლიოგრაფიულ სიაში მიუთითეთ ჯერ სამამულო, შემდეგ უცხოელი ავტორები (გვარი, ინიციალები, სტატიის სათაური, ჟურნალის დასახელება, გამოცემის ადგილი, წელი, ჟურნალის №, პირველი და ბოლო გვერდები). მონოგრაფიის შემთხვევაში მიუთითეთ გამოცემის წელი, ადგილი და გვერდების საერთო რაოდენობა. ტექსტში კვადრატულ ფხიხლებში უნდა მიუთითოთ ავტორის შესაბამისი N ლიტერატურის სიის მიხედვით. მიზანშეწონილია, რომ ციტირებული წყაროების უმეტესი ნაწილი იყოს 5-6 წლის სიღრმის.

9. სტატიას თან უნდა ახლდეს: ა) დაწესებულების ან სამეცნიერო ხელმძღვანელის წარდგინება, დამოწმებული ხელმოწერითა და ბეჭდით; ბ) დარგის სპეციალისტის დამოწმებული რეცენზია, რომელშიც მითითებული იქნება საკითხის აქტუალობა, მასალის საკმაობა, მეთოდის სანდოობა, შედეგების სამეცნიერო-პრაქტიკული მნიშვნელობა.

10. სტატიის ბოლოს საჭიროა ყველა ავტორის ხელმოწერა, რომელთა რაოდენობა არ უნდა აღემატებოდეს 5-ს.

11. რედაქცია იტოვებს უფლებას შეასწოროს სტატია. ტექსტზე მუშაობა და შეჯერება ხდება საავტორო ორიგინალის მიხედვით.

12. დაუშვებელია რედაქციაში ისეთი სტატიის წარდგენა, რომელიც დასაბეჭდად წარდგენილი იყო სხვა რედაქციაში ან გამოქვეყნებული იყო სხვა გამოცემებში.

აღნიშნული წესების დარღვევის შემთხვევაში სტატიები არ განიხილება.

| | |
|--|---------|
| Gavrilova Uliana V, Alieva Samaya A, Gerasimenko Aleksandra A, Mikhaleva Ekaterina A, Solovieva Ekaterina V, Vedzizheva Khava Kh, Sadykov Magomed L, Belousova Anastasia A, Ladaev Abubakar Kh, Aupov Ibragim A, Maasheva Elita A, Makhamaev Ilias B, Yanarbaev Ibragim R. NEXT-GENERATION ANTIOXIDANTS: SHOULD WE TARGET PEROXIREDOXINS (PRX)?..... | 6-14 |
| Marina Endeladze, Maia Zhamutashvili, Tinatin Gognadze, Elene Meskhi, Natia Jojua, M. Akhvlediani. CASE REPORT OF CAT SCRATCH DISEASE (BARTONELLA)..... | 15-17 |
| Karapetyan A.G, Santini C, Pellei M, Caviglia M, Dallakyan A.M, Petrosyan Zh.H, Danielyan M.H, Nebogova K.A, Grigoryan V.S. MANIFESTATION OF RADIOPROTECTIVE PROPERTIES IN COPPER COMPLEXES [CU(L ^{CE3}) ₂] AND [CU(ADM)(PPH ₃) ₂]PF ₆ | 18-22 |
| Nato Nakudashvili, Levan Ratiani, Tamar Megrelishvili, Elene Saribekovi, Marine Tsabadze, Nina Kipiani, Nino Intskirveli, Magda Tortladze, Tea Gabunia, Shorena Tsiklauri, Zaza Nakudashvili, George Ormotsadzr, Tamar Sanikidze. FEATURES OF VASOMOTOR RHINITIS (VMR) IN PATIENTS WITH A HISTORY OF COVID-19 INFECTION..... | 23-31 |
| Warda Mohamed, Rashid Eltayeb, Hussam Ali Osman, Mosab Omer, Elryah. I. Ali, Ammar Abdelmola, Tagwa Yousif, Mohamed Belhocine, Safa Taha, Elyasa Elfaki, Wael Alzahrani, Asaad Babker, Abdelgadir Elamin Eltom, Marwan Ismail, Ayman Alfeel. BIOCHEMICAL INSIGHTS AND HORMONAL MARKERS OF POLYCYSTIC OVARY SYNDROME IN SUDANESE WOMEN: EXPLORING INFERTILITY AND RISK FACTORS IN RED SEA STATE..... | 32-39 |
| Kolupayev S.M, Goloborodko M.M, Bytiak S.Yu, Lavrinenko A.S, Lupyr M, Lantukh I.V, Lytvynova I.L, Gulbs O.A, Dikhtyarenko S.Yu, Kobets O.V. PSYCHOLOGICAL FEATURES OF THE REHABILITATION OF PERSONS WITH UROLITHIASIS..... | 40-44 |
| Wang-Yong Li, Xiao-Yan Yang, Zhun Cai, Guo-Fu Chen. A CASE OF CECAL CANCER WITH RETROPERITONEAL ABSCESS..... | 45-48 |
| Tamriko Dzotsenidze, Arsen Gvenetadze, Giorgi Burkadze, Ekaterine Isakidi, Mariam Shulaia. GENETIC ALTERATIONS IN TUBO-OVARIAN EPITHELIUM DURING OVARIAN NEOPLASIA..... | 49-54 |
| Ali M. Muhammed Ali, Omar M. Yahya, Ehsan HT. AlDabbagh. IN SILICO DOCKING OF SILYMARIN ACTIVE CONSTITUENTS WITH INSULIN RECEPTORS: A STEP TOWARD DIABETES THERAPEUTICS..... | 55-62 |
| Armenuhi Avagyan, Taline K. Ashekian, Armenuhi Snkhchyan, Hasmik Mkrtychyan, Tigran Petrosyan. BARRIERS AND SOLUTIONS IN THE USE OF ALTERNATIVE AND AUGMENTATIVE COMMUNICATION: A PILOT DESCRIPTIVE STUDY PRESENTING INSIGHTS FROM ARMENIAN PROFESSIONALS..... | 63-71 |
| Xin-Juan Wang, Lian-Ping He. STATISTICAL TEACHING ON BUILDING STROKE PREDICTION MODELS..... | 72-75 |
| Kosherova Bakhyt Nurgaliyevna, Abbozova Shakhnosa Maratovna, Smagul Manar Asyrovna, Zhumagaliyeva Galina Dautovna, Sagyndykova Togzhan Baibolsynovna. MEASLES IN PREGNANCY IN THE REPUBLIC OF KAZAKHSTAN: CLINICAL AND LABORATORY MANIFESTATIONS AND OUTCOMES..... | 76-80 |
| Warda Mohamed, Rashid Eltayeb, Hussam Ali Osman, Mosab Omer, Elryah. I. Ali, Ammar Abdelmola, Tagwa Yousif, Mohamed Belhocine, Safa Taha, Elyasa Elfaki, Wael Alzahrani, Ayman Alfeel, Asaad Babker, Abdelgadir Elamin Eltom, Marwan Ismail. FOLLICLE-STIMULATING HORMONE RECEPTOR MUTATIONS IN SUDANESE WOMEN: A STUDY ON POLYCYSTIC OVARY SYNDROME..... | 81-86 |
| Saidulaev M.A, Osipova N.I, Gurtskaya A.D, Semov N.D, Khusainov R.R, Khabarov E.A, Lech D.S, Stotland P.A, Eloeva V.V, Syromyatnikova A.V. PIRACETAM HELPS RESTORE VISION AFTER CRANIOCEREBRAL TRAUMA..... | 87-88 |
| Ming Li, Wen-Wen Hao, Li-Juan Ru. THE ASSOCIATION BETWEEN AQUEOUS HUMOR MICROENVIRONMENT IN DIABETIC CATARACT PATIENTS AND POSTOPERATIVE MACULAR EDEMA AND VISUAL ACUITY CHANGES..... | 89-90 |
| Voloshyn-Haponov I.K, Lantukh I.V, Gulbs O.A, Dikhtyarenko S.Yu, Kobets O.V, Pustova N.O, Popova N.G, Gridneva O.V, Ostapenko V.M, Mikhhanovska N.G, Torianyik I.I. PSYCHOLOGICAL FEATURES OF THE SUBJECTIVE PERCEPTION OF THE QUALITY OF LIFE OF PATIENTS WITH HEPATOCEREBRALDYSTROPHY..... | 91-95 |
| Bangqiang Hou, Wei Liu, Ke Pan, Yiya Wang, Yaomin Luo, Yutong Han, Jingjing Liu, Qing Wu, Yinxiu Wang. BRAIN NETWORK FUNCTIONAL CONNECTIVITY AND CORTICAL ACTIVATION FEATURES DURING THE SWALLOWING TASK FOR THE PATIENTS OF POST STROKE DYSPHAGIA: A MULTI- CHANNEL FNIRS STUDY..... | 96-107 |
| Assel Ibrayeva, Dinara Ospanova, Korlan Saduakasova, Anar Akshalova, Anar Muratbayeva, Shynar Tanabayeva, Ildar Fakhradiyev. PREVALENCE AND SOCIO-DEMOGRAPHIC RISK FACTORS OF EMOTIONAL BURNOUT AMONG PSYCHIATRISTS AND NURSING STAFF IN PSYCHIATRIC SERVICES IN KAZAKHSTAN..... | 108-115 |

| | |
|---|---------|
| Violeta Grajčevci Uka, Art Uka, Lirim Isufi. THE SOCIODEMOGRAPHICAL AND MORPHOLOGICAL CHARACTERISTICS OF PRESCHOOL CHILDREN WITH SIDEROPENIC ANEMIA IN THE KOSOVO SAMPLE..... | 116-119 |
| Li-Bo Wang, Chun-Miao Xu. ENHANCING OPHTHALMIC NURSING EDUCATION: A COMPREHENSIVE APPROACH TO CLINICAL TEACHING AND TRAINING..... | 120-122 |
| Maia Zhamutashvili, Tinatin Gognadze, Natia Jojua, Elene Meskhi, Ketevan Meskhi, Ekaterine Dolmazashvili. CO-OCCURRENCE OF HANTAVIRUS PULMONARY SYNDROME AND HEMORRHAGIC FEVER WITH RENAL SYNDROME..... | 123-125 |
| Abdulkalikova D.B, Auezova A.M, Baymuratova M.A, Yessembayeva S.S, Yermukhanova G.T, Yerkibayeva ZH.U. WORKSHOP AS A PRACTICE-ORIENTED METHOD OF TEACHING DENTISTS: INTRODUCTION TO EDUCATIONAL PROGRAMS FOR WORKING WITH CHILDREN WITH AUTISM SPECTRUM DISORDERS (ASD)..... | 126-132 |
| Li-Juan Ru, Qian-Qian Yao, Ming Li. RISK PREDICTION MODEL FOR ACUTE KIDNEY INJURY IN PATIENTS WITH SEVERE ACUTE PANCREATITIS..... | 133-135 |
| Sawer S. Ahmed, Abdulazeez M. Brifkani, Haval J. Ali, Nasir A. Al Allawi. FACTORS AFFECTING HEALTH RELATED QUALITY OF LIFE IN ADULT PATIENTS WITH BETA-THALASSEMIA MAJOR..... | 136-142 |
| Liu-Xia Shi, Xiao-Ya Peng, Xiao-Xu Ruan, Rui Li, Wen-Jie Wen, Chao Deng. WHOLE TRANSCRIPTOME SEQUENCING AND CIRC_HSA_0001847 ON PROLIFERATION AND INVASION OF ORAL SQUAMOUS CELL CARCINOMA..... | 143-155 |
| Takuma Hayashi, Krishna Prasad Acharya, Sarita Phuyal, Ikuo Konishi. THE ROLE OF LIVE BIRD MARKETS SHOULD BE EMPHASIZED IN PREVENTING THE SPREAD OF HIGHLY PATHOGENIC AVIAN INFLUENZA INFECTIONS..... | 156-158 |
| Karapetyan A.G, Grigoryan V.S, Santini C, Pellei M, Del Gobbo J, Dallakyan A.M, Petrosyan Zh.H, Fanarjyan S.A, Danielyan M.H, Nebogova K.A. BLOOD AND CYTOGENETIC MARKERS IN EXPERIMENTAL BURNS AND THEIR TREATMENT WITH [CU(L ^{CF3}) ₂] AND [CU(ADM)(PPH ₃) ₂]PF ₆ | 159-163 |
| Sharofova M.U, Khalimova F.T, Habasi Maidina, Jiangyu Zhao, Haji Akber Aisa. ANTIMICROBIAL AND ANTI-INFLAMMATORY ACTIVITY OF PLANT EXTRACTS: PROSPECTS FOR THE DEVELOPMENT OF COMBINED THERAPEUTIC AGENTS..... | 164-168 |

THE ROLE OF LIVE BIRD MARKETS SHOULD BE EMPHASIZED IN PREVENTING THE SPREAD OF HIGHLY PATHOGENIC AVIAN INFLUENZA INFECTIONS

Takuma Hayashi^{1*}, Krishna Prasad Acharya², Sarita Phuyal³, Ikuo Konishi¹.

¹National Hospital Organization, Kyoto Medical Center, Cancer Medicine, Kyoto, Kyoto, Japan.

²Animal Disease Investigation and Control Division (ADICD), Department of Livestock Services (DLS), Hariharbhawan, Lalitpur, Nepal.

³Central Referral Veterinary Hospital (CRVH), Department of Livestock Services (DLS), Tripureshwar, Kathmandu, Nepal.

Abstract.

Zoonotic diseases refer to infectious diseases caused by pathogens that infect or parasitize both humans and other vertebrate animals. In recent years, the novel coronavirus disease (COVID-19) has been known to be a zoonotic disease. Academic fields related to zoonotic diseases include veterinary medicine and virology. According to a report by the Food and Agriculture Organization of the United Nations (FAO), approximately 70% of new human infectious diseases identified since 1940 are of animal origin. On the other hand, approximately twice as many pathogens are transmitted from humans to animals as are transmitted from animals to humans. In other words, humans are a major source of infection for animals. Due to increased opportunities for various animals to be imported and kept as pets, pathogens that were previously rare or unknown can suddenly appear in human society. Alternatively, there are ethnic groups in the world who eat wild birds and animals that are not sufficiently cooked, and there have been cases of pathogens infecting wild birds and animals being transmitted to humans. If a zoonotic disease caused by a new pathogen like this becomes prevalent as an emerging infectious disease, there is a high risk of it becoming a pandemic because humans have not yet acquired immunity to the new pathogen. This problem was clearly evident in the Severe Acute Respiratory Syndrome (SARS) pandemic that emerged in 2003, and the COVID-19 pandemic caused by infection with SARS-CoV-2, which was identified in 2019. In this review, we discuss the importance of management practices and current information on zoonotic diseases in outlets selling live wild birds and wildlife for consumption.

Key words. Highly pathogenic avian influenza, H5N1, influenza virus, Bird flu.

Introduction.

Highly pathogenic avian influenza (HPAI) is a viral disease of birds with zoonotic potential caused by influenza virus belonging to the family orthomyxoviridae. It is one of the most economically important diseases of birds causing a huge economic burden to poultry sectors worldwide and it is endemic in many countries in Asia and Africa [1-4]. Thus, great strides have been made by different countries to prevent and control HPAI. However, they seem to have failed to effectively control HPAI, which is partly attributed to the fact that control programs have failed to include some of the key drivers of HPAI transmission such live bird markets (LBMs).

The H5N1 virus that infects dairy cows genetically modifies in the living dairy cow, increasing its infectivity to humans.

LBMs are an essential component of poultry marketing systems in low- and middle-income countries, where people

prefer to purchase live poultry for consumption rather than frozen meat. These LBMs, where birds of different species are brought together from different geographical areas, facilitate the mixed selling of live poultry birds which in turn facilitate the transmission, evolution, and spillover of pathogens carried by marketed birds [5]. Furthermore, LBMs create a network of contacts between poultry birds and the human, which may pose a danger to public health by facilitating the transmission of zoonotic diseases if proper hygiene sanitation and biosecurity measures are not followed. Several previous studies have found that LBMs are hotspots for avian influenza viruses [6-12] and are linked to outbreaks of HPAI [5,6,13-16]. For instance, the human outbreak of H5N1 in 1997 [13,14] and the SARS outbreak in 2003 [12,17] were both linked to LBMs. Furthermore, these LBMs are found to have poor hygienic practices, poor biosecurity measures and a high density of birds and humans, all of which facilitate the transmission HPAI virus [6,12,18-21]. Furthermore, the frequent introduction of susceptible, unimmunized bird populations increases the numbers of susceptible animal populations in LBMs as well as the likelihood of ongoing transmission of viruses to other areas. Besides that, the frequent movement of humans to and from LBMs contributes to their spread to humans. This was the case with H5N1, where the strain was developed through genetic re-assortment in poultry birds and got spillover into humans [22]. Given the close contact, they may spread into susceptible bird populations and humans, entering either the poultry value chain or food chain systems, regardless of how stringent the surveillance systems and biosecurity measures are. Thus, there is a high chance that those live bird markets are helping the spread of the HPAI virus. Furthermore, there could be circulation of more than one strain of virus whose genome may recombine and lead to the development of new novel variants that are more virulent as already evinced in the case of the H5N1 [22,23]. Specifically, when H5N1, which infects farmed poultry, infects dairy cows, genetic modification occurs within the body of the cow. As a result, it has been shown that a strain of H5N1 that is highly infectious to humans is produced (Figure 1). In recent years, there have also been reports of H5N8 [24-27] and H7N9 [25,28] strains developing through genetic re-assortment and spillover into human population, which could later may turn into a global pandemic by acquiring sustained human-human transmissibility.

Hygiene management of LBM is important to prevent the spread of zoonotic diseases.

Given their role avian influenza transmission dynamics, including that of HPAI, LBMs have been a primary target for AIV control strategies aimed at reducing virus circulation in the LBMs. For example, during a zoonotic outbreak of H7N9




Figure 1. Genetic reassortment.

When a given cell is infected by two H5N1 viral strains, hybrid viruses can be generated by exchanging their segmented genome, a mechanism called “genetic reassortment. Dairy cows can be infected by both avian flu and human influenza viruses, because the glycans in the upper respiratory tract express sialic acids with not only an α -2,3 linkage but also an α -2,6 linkage. When avian and human influenza viruses coinfect dairy cows, hybrid viruses could be generated in the body of dairy cows. The infectivity of the hybrid viruses appears to be enhanced.

in China in 2013, closing LBMs was shown to be remarkably effective in reducing the risk of human infection by up to 99% [13,15]. Control strategies implemented in LBMs have also been found to significantly reduce AIV detection rates in poultry [13]. A monthly rest days with a routine market closure followed by the slaughter of unsold poultry, a ban on the sale of live quail, and a ban on overnight storage of live poultry all helped to control HPAI in Hong Kong [16]. Therefore, it is clear that LBMs may help in the maintenance and transmission of the HPAI virus, and their role in the transmission of the HPAI virus should not be overlooked.

It is necessary to conduct research to understand the transmission dynamics of HPAI in LBMs while taking into account the local ecological and epidemiological drivers as well as to assess the risk of potential maintenance and transmission to poultry birds and spillover into humans leading to public health risks. It is imperative to develop and implement stringent and meticulous surveillance systems based on “rigorous scientific risk assessment” to detect viruses on time, as well as devise and implement effective preventive and control strategies based on “rigorous scientific risk assessment”, improve the hygiene and sanitation of the LBMs, and prevent mixing of birds of different species, to prevent the transmission and spillover of viruses so as to mitigate the threat of LBMs acting as reservoirs or sources of HPAI and safeguard both the associated animal and public health.

Competing interests:

The authors declare no conflict of interest.

Author contribution:

TH, KPA, and SP were involved in the study design, data collection, data review and interpretation, and manuscript writing. TH, KPA, and SP were involved in the literature search, study design, data collection, data interpretation, and manuscript writing. TH, KPA, and SP were involved in data collection and interpretation. TH, KPA, and SP were involved

in data collection and interpretation and manuscript writing. TH, KPA, and SP were involved in the study conception and design, data analysis and interpretation, and manuscript writing. TH and IK were the medical leads for AstraZeneca, and they participated in the data collection and evaluation and manuscript writing and editing. TH and IK were the lead physicians and were involved in the study design and conduct, data analysis and interpretation, and manuscript review.

Funding:

This clinical research was performed using research funding from the following: Japan Society for Promoting Science for TH (grant no. 19K09840), START-program Japan Science and Technology Agency (JST) for TH (grant no. STSC20001), National Hospital Organization Multicenter Clinical Study for TH (grant no. 2019-Cancer in general-02), and Japan Agency for Medical Research and Development (AMED) (grant no. 22ym0126802j0001), Tokyo, Japan. The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Data Availability Statement:

Data are available on various websites and have also been made publicly available. More information can be found in the first paragraph of the Results section. The transparency document associated with this article can be found in the online version at https://kyoto.hosp.go.jp/html/guide/medicalinfo/clinical_research/expand/gan.html (accessed on 18 March 2024).

Informed Consent Statement:

This research is not clinical research, therefore Informed consent is not required.

Acknowledgments:

The authors want to thank Dr. Yoshihiro Kawaoka at the institution of medical science, The University of Tokyo for providing clinical research information. The authors also want to acknowledge all medical staff for clinical research at Kyoto

University School of Medicine and the National Hospital Organization Kyoto Medical Center.

REFERENCES

1. Chowdhury S, Hossain M.E, Ghosh P.K, et al. The pattern of highly pathogenic avian influenza H5N1 outbreaks in South Asia. *Trop. Med. Infect. Dis.* 2019;4:138.
2. Durand L.O, Glew P, Gross D, et al. Timing of influenza a(H5n1) in poultry and humans and seasonal influenza activity worldwide, 2004–2013. *Emerg. Infect. Dis.* 2015;21:202-208.
3. Pfeiffer D.U, Otte M.J, Roland-Holst D, et al. A one health perspective on HPAI H5N1 in the Greater Mekong sub-region. *Comp. Immunol. Microbiol. Infect. Dis.* 2013;36:309-319.
4. Letsholo S.L, James J, Meyer S.M, et al. Emergence of High Pathogenicity Avian Influenza Virus H5N1 Clade 2.3.4.4b in Wild Birds and Poultry in Botswana. *Viruses* 2022;14:2601.
5. Moyen N, Hoque M.A, Mahmud R, et al. Avian influenza transmission risk along live poultry trading networks in Bangladesh. *Sci. Rep.* 2021;11.
6. Fournié G, Guitian J, Desvaux S, et al. Identifying live bird markets with the potential to act as reservoirs of avian influenza a (H5N1) virus: A survey in Northern Viet Nam and Cambodia. *PLoS One.* 2012;7.
7. Indriani R, Samaan G, Gultom A, et al. Environmental sampling for avian influenza virus a (H5N1) in live-bird markets, Indonesia. *Emerg. Infect. Dis.* 2010;16:1889-1895.
8. Coker R.J, Hunter B.M, Rudge J.W, et al. Emerging infectious diseases in southeast Asia: Regional challenges to control. *Lancet.* 2011;377:599-609.
9. Rimi N.A, Hassan M.Z, Chowdhury S, et al. A decade of avian influenza in Bangladesh: Where are we now?, *Trop. Med. Infect. Dis.* 2019;4.
10. Islam A, Rahman M.Z, Hassan M.M, et al. Determinants for the presence of avian influenza virus in live bird markets in Bangladesh: Towards an easy fix of a looming one health issue. *One Heal.* 2023;17.
11. Pinotti F, Kohnle L, Lourenço J, et al. Modelling the transmission dynamics of H9N2 avian influenza viruses in a live bird market. *Nat. Commun.* 2024;15:1-10.
12. Webster R.G. Wet markets - A continuing source of severe acute respiratory syndrome and influenza?. *Lancet.* 2004;363:234-236.
13. Gao G.F. Influenza and the live poultry trade. *Science.* 2014;344:235.
14. Wan X.-F, Dong L, Lan Y, et al. Indications that Live Poultry Markets Are a Major Source of Human H5N1 Influenza Virus Infection in China. *J. Virol.* 2011;85:13432-13438.
15. Yu H, Wu J.T, Cowling B.J, et al. Effect of closure of live poultry markets on poultry-to-person transmission of avian influenza A H7N9 virus: An ecological study. *Lancet.* 2014;383:541-548.
16. Connie Leung Y.H, Lau E.H.Y, Zhang L.J, et al. Avian influenza and ban on overnight poultry storage in live poultry markets, Hong Kong. *Emerg. Infect. Dis.* 2012;18:1339-1341.
17. Guan Y, Zheng B.J, He Y.Q, et al. Isolation and characterization of viruses related to the SARS coronavirus from animals in Southern China. *Science.* 2003;302:276-278.
18. Fournié G, Guitian F.J, Mangtani P, et al. Impact of the implementation of rest days in live bird markets on the dynamics of H5N1 highly pathogenic avian influenza. *J. R. Soc. Interface.* 2011;8:1079-1089.
19. Turner J.C.M, Feeroz M.M, Hasan M.K, et al. Insight into live bird markets of Bangladesh: An overview of the dynamics of transmission of H5N1 and H9N2 avian influenza viruses. *Emerg. Microbes Infect.* 2017;6:e12.
20. Negovetich N.J, Feeroz M.M, Jones-Engel L, et al. Live bird markets of bangladesh: H9n2 viruses and the near absence of highly pathogenic h5n1 influenza. *PLoS One.* 2011;6.
21. Biswas P.K, Giasuddin M, Nath B.K, et al. Biosecurity and Circulation of Influenza A (H5N1) Virus in Live-Bird Markets in Bangladesh, 2012, *Transbound. Emerg. Dis.* 2017;64:883-891.
22. Xu X, Subbarao K, Cox N.J, et al. Genetic characterization of the pathogenic influenza A/Goose/Guangdong/1/96 (H5N1) virus: Similarity of its hemagglutinin gene to those of H5N1 viruses from the 1997 outbreaks in Hong Kong. *Virology.* 1999;261:15-19.
23. Guan Y, Peiris M, Kong K.F, et al. H5N1 influenza viruses isolated from geese in southeastern China: Evidence for genetic reassortment and interspecies transmission to ducks. *Virology.* 2002;292:16-23.
24. Kang H.M, Lee E.K, Song B.M, et al. Novel reassortant influenza a(H5N8) viruses among inoculated domestic and wild ducks, South Korea, 2014. *Emerg. Infect. Dis.* 2015;21:298-304.
25. Huang P, Sun L, Li J, et al. Potential cross-species transmission of highly pathogenic avian influenza H5 subtype (HPAI H5) viruses to humans calls for the development of H5-specific and universal influenza vaccines. *Cell Discov.* 2023;58.
26. Jeong J, Kang H.M, Lee E.K, et al. Highly pathogenic avian influenza virus (H5N8) in domestic poultry and its relationship with migratory birds in South Korea during 2014. *Vet. Microbiol.* 2014;173:249-257.
27. ECDC. First identification of human cases of avian influenza A (H5N8) infection. 2021. <https://www.ecdc.europa.eu/sites/default/files/documents/First-identification-human-cases-avian-influenza-A-H5N8-infection.pdf>.
28. Chen J, Zhang J, Zhu W, et al. First genome report and analysis of chicken H7N9 influenza viruses with poly-basic amino acids insertion in the hemagglutinin cleavage site. *Sci. Rep.* 2017;7.